

Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11) **EP 0 506 477 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention  
of the grant of the patent:  
23.06.1999 Bulletin 1999/25

(51) Int. Cl.<sup>6</sup>: **C07K 14/575**, C12N 15/19,  
C12N 5/10, A61K 38/22  
// C07H21/00

(21) Application number: **92302750.2**

(22) Date of filing: **27.03.1992**

(54) **Vascular endothelial cell growth factor C subunit**

Untereinheit-C des vaskulären endothelialen Zellwachstumsfaktors

Sous-unité C du facteur de croissance de cellules vasculaires endothéliales

(84) Designated Contracting States:  
**CH DE FR GB IT LI NL**

(30) Priority: **28.03.1991 US 676436**

(43) Date of publication of application:  
**30.09.1992 Bulletin 1992/40**

(73) Proprietor: **Merck & Co., Inc.**  
**Rahway New Jersey 07065-0900 (US)**

(72) Inventors:  
• **Bayne, Marvin L.**  
**Westfield, NJ 07090 (US)**  
• **Thomas, Kenneth A., Jr.**  
**Chatham Burrough, NJ 07928 (US)**

(74) Representative:  
**Thompson, John Dr. et al**  
**Merck & Co., Inc.**  
**European Patent Department**  
**Terlings Park**  
**Eastwick Road**  
**Harlow, Essex CM20 2QR (GB)**

(56) References cited:  
**EP-A- 0 370 989**                      **EP-A- 0 399 816**  
**EP-A- 0 476 983**                      **WO-A-90/13649**  
**WO-A-91/02058**                      **WO-A-92/06194**  
  
• **PROCEEDINGS OF THE NATIONAL ACADEMY**  
**OF SCIENCES OF USA. vol. 88, October 1991,**  
**WASHINGTON US pages 9267 - 9271; D.**  
**MAGLIONE ET AL.: 'Isolation of a human**  
**placenta cDNA coding for a protein related to the**  
**vascular permeability factor'**

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

**EP 0 506 477 B1**

**Description****BRIEF DESCRIPTION OF THE DRAWING****[0001]**

Figure 1. Full length amino acid residue protein translation product and its cDNA coding sequence for VEGF AA subunit A plus polypeptide cleavage products used to determine the amino acid sequence.

Figure 2. Full length amino acid residue protein translation product and its cDNA coding sequence for VEGF AB subunit A plus polypeptide cleavage products used to determine the amino acid sequence.

Figure 3. Full length amino acid residue protein translation product and its cDNA coding sequence for VEGF AB subunit B plus polypeptide cleavage products used to determine the amino acid sequence.

Figure 4. Full cDNA coding sequence and full length amino acid residue protein translation product for VEGF A 146 amino acid residue subunit SEQ ID NOS:23 & 33.

Figure 5. Full cDNA coding sequence and full length amino acid residue protein translation product for VEGF A 190 amino acid residue subunit SEQ ID NOS:30 & 31.

Figure 6. Full cDNA coding sequence and full length amino acid residue protein translation product for VEGF A 214 amino acid residue subunit SEQ ID NOS:34 & 35.

Figure 7. Full cDNA coding sequence and full length amino acid residue protein translation product for VEGF B 138 amino acid residue subunit SEQ ID NOS:36 & 37.

Figure 8. Full cDNA coding sequence and full length amino acid residue protein translation product for VEGF B 158 amino acid residue subunit SEQ ID NOS:38 & 39.

Figure 9. Full cDNA coding sequence and full length amino acid residue protein translation product for VEGF C 154 amino acid residue subunit SEQ ID NOS:40 & 41.

**BACKGROUND OF THE INVENTION**

[0002] A new class of cell-derived dimeric mitogens with apparently restricted specificity for vascular endothelial cells has recently been identified and generally designated vascular endothelial growth factors (VEGFs). The mitogen has been purified from: conditioned growth media of rat glioma cells, [Conn *et al.*, Proc. Natl. Acad. Sci. USA **87**: 1323-1327 (1990)]; conditioned growth media of bovine pituitary folliculo stellate cells [Ferrara and Henzel, Biochem. Biophys. Res. Comm. **161**: 851-858 (1989) and Gospodarowicz *et al.*, Proc. Natl. Acad. Sci. USA **86**: 7311-7315 (1989)]. An endothelial cell growth factor isolated from mouse neuroblastoma cell line NB41 with an unreduced molecular mass of 43-51 kDa and a reduced mass of 23-29 kDa has been described by Levy *et al.*, Growth Factors **2**: 9-19 (1989). Connolly *et al.* (J. Biol. Chem. **264**: 20017-20024 [1989]; J. Clin. Invest. **84**: 1470-1478 [1989]) describe a human vascular permeability factor that stimulates vascular endothelial cells to divide *in vitro* and promotes the growth of new blood vessels when administered into healing rabbit bone grafts or rat corneas. An endothelial cell growth factor has been purified from the conditioned medium of the AtT-20 pituitary cell line by Plouet *et al.*, EMBO Journal **8**: 3801-3806 (1989). The growth factor was characterized as a heterodimer composed of subunits with molecular mass of 23 kDa. Leung *et al.* (Science **246**: 1306-1309 [1989]), Keck *et al.* (Science **246**: 1309-1312 [1989]) and Conn *et al.* (Proc. Natl. Acad. Sci. USA **87**: 2628-2632 [1990]) have described cDNAs which encode VEGF A which is homologous to the A and B chains of platelet-derived growth factor. Vascular endothelial growth factor I (VEGF I, VEGF AA) is a homodimer with an apparent molecular mass of 46 kDa, with each subunit having an apparent molecular mass of 23 kDa. VEGF I has distinct structural similarities to platelet-derived growth factor (PDGF), a mitogen for connective tissue cells but not vascular endothelial cells from large vessels.

**OBJECTS OF THE INVENTION**

[0003] It is, accordingly, an object of the present invention to provide novel vascular endothelial growth factor C subunit DNA free of other mammalian DNA. Another object is to provide recombinant genes capable of expressing VEGF C subunit monomer or dimer. Another object is to provide vectors containing the DNA sequences for VEGF A or B plus C subunits. A further object is to provide a host cell transformed with a vector containing the DNA sequence for VEGF A or B plus C or VEGF C alone. It is also an object to provide a recombinant process for making VEGF C subunit. Another object is to provide a novel vascular endothelial cell growth factor which contains the C subunit. This may include heterodimers AC and BC and homodimer CC.

SUMMARY OF THE INVENTION

[0004] Vascular endothelial cell growth factor C subunit DNA is prepared by polymerase chain reaction techniques. The DNA encodes a protein that may exist as either a heterodimer or homodimer. The protein is a mammalian vascular endothelial cell mitogen and as such is useful for the promotion of vascular development and repair. This unique growth factor is also useful in the promotion of tissue repair.

DETAILED DESCRIPTION

[0005] The present invention relates to a unique vascular endothelial cell growth factor (designated VEGF), isolated and purified from glioma cell conditioned medium, which exhibits mitogenic stimulation of vascular endothelial cells. Glioma is defined herein as any neoplasm derived from one of the various types of cells that form the interstitial tissue of the central nervous system including brain, spinal cord, posterior pituitary gland and retina. Consequently, the scope of the present invention is intended to include the unique growth factor isolated and purified from any mammalian tissue or other cells including cell lines. Cell lines include, but are not limited to, glioma-derived cell lines such as C6, hs 683 and GS-9L; glioblastomas such as A-172 and T98G; neuroblastomas such as IMR-32 and SK-N-MC; neurogliomas such as H4; tetromas such as XB-2; astrocytomas such as U-87 MG and U-373 MG; embryonal carcinomas and non-transformed glial or astrocyte cell lines, and the human medulloblastoma line TE 671, with GS-9L and TE 671 being preferred. VEGF AB is present and can be isolated from rat tissue including ovary, heart and kidney. Anterior pituitary tumor cell lines such as GH3 and Hs 199 may also be used. It is intended that VEGF of this invention can be obtained from any mammal species capable of producing VEGF, this includes, but is not limited to, rat and human.

[0006] Vascular endothelial cell growth factor may exist in various microheterogeneous forms which are isolated from one or more of the various cells or tissues described above. Microheterogeneous forms as used herein refer to a single gene product, that is a peptide produced from a single gene unit of DNA, which is structurally modified at the mRNA level or following translation. Peptide and protein are used interchangeably herein. The microheterogeneous forms will all have similar mitogenic activities. Biological activity and biologically active are used interchangeably and are herein defined as the ability of VEGF to stimulate DNA synthesis in target cells including vascular endothelial cells as described below which results in cell proliferation. The modifications may take place either *in vivo* or during the isolation and purification process. *In vivo* modification results from, but is not limited to, proteolysis, glycosylation, phosphorylation, deamidation or acetylation at the N-terminus. Proteolysis may include exoproteolysis wherein one or more terminal amino acids are sequentially, enzymatically cleaved to produce microheterogeneous forms which have fewer amino acids than the original gene product. Proteolysis may also include endoproteolytic modification that results from the action of endoproteases which cleave the peptide at specific locations within the amino acid sequence. Similar modifications can occur during the purification process which also results in production of microheterogeneous forms. The most common modification occurring during purification is proteolysis which is generally held to a minimum by the use of protease inhibitors. Under most conditions one or more microheterogeneous forms are present following purification of native VEGFs. Native VEGFs refers to VEGF isolated and purified from cells that produce VEGFs. Vascular endothelial cell growth factor may also exist in various alternatively spliced forms which is defined herein as the production of related mRNAs by differential processing of exons and introns. Exons are defined as those parts of the DNA sequence of a eukaryotic gene that code for the final protein product. It is also intended that the present invention includes VEGF subunits C which are defined as comprising the full length translation products of all alternatively spliced mRNAs made from the gene encoding the subunits and their corresponding mature amino acid sequences generated by proteolytic removal of the amino terminal secretory leader amino acid sequences. It is further intended that the invention only include those microheterogeneous and alternatively spliced VEGF subunits which when in the dimeric form exhibit biological activity such as vascular endothelial cell stimulation as discussed below.

[0007] Glioma cells such as the rat cell line GS-9L are grown to confluence in tissue culture flasks, about 175 cm<sup>2</sup>, in a cell culture medium such as Dulbecco's Modified Eagle's Medium (DMEM) supplemented with about 10% newborn calf serum (NCS). When the cells reach confluence the culture medium is removed, the cell layers are washed with Ca<sup>++</sup>, Mg<sup>++</sup>-free phosphate buffered saline (PBS) and are removed from the flasks by treatment with a solution of trypsin, about 0.1%, and EDTA, about 0.04%. The cells, about 1 x 10<sup>8</sup>, are pelleted by centrifugation, resuspended in about 1500 ml of DMEM containing about 5% NCS and plated into a ten level cell factory (NUNC), 6,000 cm<sup>2</sup> surface area. The cells are incubated for about 48 to about 96 hours, with 72 hours preferred, at about 37° C in an atmosphere of about 5% CO<sub>2</sub>. Following incubation the medium is removed and the cell factories are washed about 3 times with PBS. About 1500 ml of fresh culture media is added containing about a 1:2 mixture of Ham's-F12/DMEM containing about 15 mM Hepes, pH about 7.4, about 5 µg/ml insulin, about 10 µg/ml transferrin and with or without about 1.0 mg/ml bovine serum albumin. This medium is replaced with fresh medium after about 24 hr and collected every 48 hr thereafter. The collected conditioned medium is filtered through Whatmen #1 paper to remove cell debris and stored at about -20° C.

[0008] The GS-9L conditioned medium is thawed and brought to pH 6.0 with 1 M HCl. The initial purification step consists of cation exchange chromatography using a variety of cation exchangers on a variety of matrices such as CM Sephadex C-50, Pharmacia Mono S, Zetachrom SP and Polyaspartic Acid WCX (Nest Group) with CM Sephadex C-50 (Pharmacia) being preferred. The VEGF-containing culture medium is mixed with CM Sephadex C-50 at about 2 gm per about 20 L of the conditioned medium and stirred at low speed for about 24 hr at 4° C. The resin is allowed to settle and the excess liquid is removed. The resin slurry is packed into a column and the remaining culture medium is removed. Unbound protein is washed from the column with 0.05 M sodium phosphate, about pH 6.0, containing 0.15 M NaCl. The VEGF AB is eluted with about 0.05 M sodium phosphate, about pH 6.0, containing about 0.6 M NaCl.

[0009] The active fractions collected from the CM Sephadex C-50 column are further fractionated by lectin affinity chromatography for additional purification of VEGF AB. The lectins which may bind VEGF AB include, but are not limited to, lectins which specifically bind mannose residues such as concanavalin A and lens culinaris agglutinin, lectins which bind N-acetylglucosamine such as wheat germ agglutinin, lectins that bind galactose or galactosamine and lectins which bind sialic acids, with concanavalin A (Con A) being preferred. A 0.9 cm diameter column containing about 5 ml packed volume of Con A agarose (Vector Laboratories) is washed and equilibrated with about 0.05 M sodium acetate, about pH 6.0, containing about 1 mM  $\text{CaCl}_2$ , about 1 mM  $\text{MnCl}_2$  and about 0.6 M NaCl. The unbound protein is washed from the column with equilibration buffer. The VEGF AB is eluted with about 0.1 M NaCl buffer containing about 0.32 M  $\alpha$ -methyl mannoside and about 0.28 M  $\alpha$ -methyl glucoside.

[0010] The VEGF AB active eluate from the Con-A column is applied to a Polyaspartic Acid WCX cation exchange high performance liquid chromatography (HPLC) column, 4.6 mm x 250 mm, pre-equilibrated in about 0.05 M sodium phosphate buffer, pH 6.0. The column is eluted with a linear gradient of about 0 to 0.75 M NaCl in the phosphate buffer over about 60 minutes. The flow rate is maintained at about 0.75 ml/min collecting 0.75 ml fractions. Vascular endothelial cell growth factor AB activity is present in fractions eluting between approximately 21.7 and 28.5 ml.

[0011] The active fractions eluted from the polyaspartic WCX column that contain VEGF AB are pooled, adjusted to about pH 7.0 and loaded onto a 1 x 10 cm column of Pharmacia Chelating Sepharose 6B charged with an excess of copper chloride and equilibrated in about 0.05 M sodium phosphate, about pH 7.0, containing about 2 M NaCl and about 0.5 mM imidazole (A buffer). VEGF AB is eluted from the column with a gradient from 0-20% B over 10 minutes, 20-35% B over 45 minutes and 35-100% B over 5 minutes at a flow rate of 0.3 ml/min, where B buffer is 0.05 M sodium phosphate, pH 7.0, containing about 2 M NaCl and 100 mM imidazole. The active fractions containing VEGF AB activity eluted between about 12.6 and 22.8 ml of the gradient effluent volume.

[0012] The pooled fractions containing VEGF AB activity eluted from the metal chelate column are loaded onto a 4.6 mm x 5 cm Vydac

$\text{C}_4$  reverse phase HPLC column (5  $\mu\text{m}$  particle size) previously equilibrated in solvent A [0.1% trifluoroacetic acid (TFA)]. The column is eluted with a linear gradient of about 0 to 30% solvent B over 15 minutes, 30% B for an additional 15 minutes, then 30-45% B over 22.5 minutes and finally 45-100% B over 5.5 minutes. Solvent B consists of solvent A containing 67% acetonitrile (v/v). The flow rate is maintained at about 0.75 ml/min and fractions are collected every minute. The homogeneous VEGF AB elutes from the  $\text{C}_4$  column under these conditions at between about 32 and about 38 ml of the gradient effluent volume.

[0013] Purity of the protein is determined by sodium dodecylsulfate (SDS) polyacrylamide gel electrophoresis (PAGE) in 12.5% crosslinked gels using the technique of Laemmli, Nature 227: 680-684 (1970). The silver stained gels show VEGF AB to consist of one band under non-reducing conditions with an approximate apparent molecular mass of about 58,000 daltons. When a sample containing the microheterogeneous forms of VEGF AB is separated under reducing conditions it migrates as two about 23 kilodalton (kDa) subunits. The purification process results in VEGF AB that is essentially free of other mammalian cell products, such as proteins. Recombinantly derived VEGF AB will also be free of mammalian cell products.

[0014] Biological activity is determined by mitogenic assay using mammalian vascular endothelial cells. Human umbilical vein endothelial (HUVE) cells are plated on gelatin-coated dishes at a density of about 5000 cells per well in about 500  $\mu\text{l}$  of Medium 199 (M199) containing about 20% heat-inactivated fetal calf serum (FCS). Samples to be assayed are added at the time of plating. The tissue culture plates are incubated at about 37° C for about 12 hours and about 2 microcuries of tritiated thymidine (NEN, 20 Ci/mmol) is added per ml of assay medium (1.0  $\mu\text{Ci}/\text{well}$ ). The plates are incubated for a further 60 hr, the assay medium is removed and the plates are washed with Hanks balanced salt solution containing about 20 mM Hepes, about pH 7.5, and about 0.5 mg/ml bovine serum albumin. The cells are lysed and the labelled DNA solubilized with about 200  $\mu\text{l}$  of a solution containing about 2 gm of sodium carbonate and about 400 mg sodium hydroxide in about 100 ml water. The incorporated radioactivity was determined by liquid scintillation counting. The concentration of VEGF which elicited a half-maximal mitogenic response in HUVE cells was approximately  $2 \pm 1$  ng/ml. The glycosaminoglycan heparin, which is required in these assays at a level of 10-100  $\mu\text{g}/\text{ml}$  to promote a response to a positive control, acidic fibroblast growth factor, does not enhance mitogenic stimulation of these cells by VEGF AB.

[0015] A purified about 1-2  $\mu\text{g}$  sample of VEGF AB is reduced in about 0.1 M Tris, about pH 9.5, with about 0.1%

EDTA, about 6 M guanidinium chloride and about 20 mM dithiothreitol for about 2 hr at about 50° C. The reduced protein is carboxymethylated for about 1 hour in a solution containing about 9.2  $\mu$ M of unlabelled and 2.8  $\mu$ M of  $^{14}$ C-iodoacetic acid in about 0.7 M Tris, about pH 7.8, and about 0.1% EDTA and about 6 M guanidinium chloride. The protein is carboxymethylated for about 1 hr at room temperature. The protein is isolated after reduction and carboxymethylation by reverse phase HPLC chromatography on a Vydac C<sub>4</sub> column, about 4.6 mm x 5 cm. The protein subunits are loaded onto a column pre-equilibrated with about 0.1% TFA and eluted by a 45 ml linear gradient from about 0.1% TFA to 0.1% TFA/67% acetonitrile at a flow rate of about 0.75 ml/min. The reduced and carboxymethylated protein eluted as two peaks at approximately 23 and 25 ml with the proportion being approximately equal as determined by monitoring absorbance at 210 nm.

[0016] Samples of the reduced and carboxymethylated monomers are applied to polybrene-coated glass fiber filters and their N-terminal sequences are determined by Edman degradation in an ABI gas phase microsequencer in conjunction with an ABI 120A on line phenylthiohydantoin analyzer following the manufacturers instructions. The protein showing the peak of absorbance eluting at approximately 25 ml (A subunit or monomer) yielded an amino terminal sequence of: SEQ ID NO:1

Ala Pro Thr Thr Glu Gly Glu Gln Lys Ala His Glu Val Val which is identical to the A chain monomers of VEGF AA, Conn et al., Proc. Natl. Acad. Sci. USA 87: 2628-2632 (1990). The peak of absorbance eluting at approximately 23 ml (B subunit or monomer) yielded an N-terminal sequence of: SEQ ID NO:2

Ala Leu Ser Ala Gly Asn Xaa Ser Thr Ser Thr Glu Met Glu Val Val  
Pro Phe Asn Glu Val

plus a nearly equal amount of a truncated form of the same sequence missing the first three amino acid residues. The missing Xxx residue corresponds to an Asn residue in the cloned cDNA, see below. Since this missing Asn occurs in a classical Asn Xxx Ser/Thr N-glycosylation sequence it is presumed to be glycosylated. The A subunit and the total of both B subunits are recovered in nearly equal amounts supporting the interpretation that the two peptides combine to form an AB heterodimer in VEGF AB.

[0017] A sample of the A monomer was treated with either the protease trypsin which cleaves polypeptides on the C-terminal side of lysine and arginine residues or Lys C which cleaves polypeptides on the C-terminal side of lysine by procedures well known in the art. The peptides are isolated by reversed phase - HPLC(RP-HPLC). The amino acid sequences of the isolated peptides are determined using the Edman degradation in the ABI gas phase sequencer in conjunction with the ABI 120 A on line phenylthiohydantoin analyzer following manufacturer's instructions. The amino acid sequences are shown in Figure 1.

[0018] Reduced and carboxymethylated A monomer is dried and solubilized in about 0.7 M Tris, about pH 7.8, about 6 M guanidinium chloride containing about 0.1% EDTA. V8 protease is added in 0.1 M ammonium bicarbonate buffer, about pH 8.0, and the mixture is incubated for about 48 hr at about 37°C. The protease cleaves predominantly on the carboxyl terminal side of glutamic acid residues. The resulting polypeptides were resolved by C<sub>18</sub> RP-HPLC as above.

[0019] The reduced and carboxymethylated A subunit protein solution is adjusted to a pH of about 6.8 with 6 N HCl and dithiothreitol is added to a final concentration of 2 M for reduction of any methionine sulfoxide to methionine residues. After about 20 hr of reduction at about 39°C the protein is repurified by C<sub>4</sub> HPLC. The product is dried and cleaved on the carboxyl terminal side of methionine residues by 200  $\mu$ l of 40 mM cyanogen bromide in about 70 % (v/v) formic acid under an argon atmosphere at about 20°C for about 24 hr in the dark. The cleavage products are resolved by C<sub>18</sub> RP-HPLC. The amino acid sequence is shown in Figure 1, see Conn et al., Proc. Natl. Acad. Sci. USA 87: 2628-2632 (1990).

[0020] The full length 190 amino acid residue protein translation product of the VEGF AB, A monomer or subunit, which is now known to be identical with the VEGF AA, A monomer, and its cDNA coding sequence are shown in Figures 2 and 6. The mature amino terminus begins at residue 27, immediately following a typical hydrophobic secretory leader sequence. A single potential N-glycosylation site exists at Asn<sub>100</sub>. Most (143 amino acid residues) of the 164 residues of the reduced and carboxymethylated mature subunit including the amino terminus and HPLC reversed phase-purified products of tryptic (T), Lys-C (L), *Staphylococcus aureus* V8 protease (V8) and cyanogen bromide (CB) cleavages, were determined by direct microsequencing (Applied Biosystems 470A) using a total of 5  $\mu$ g of protein. All residues identified by amino acid sequencing are denoted by arrows pointing to the right either directly beneath the mature processed sequence following the bracket at residue 27 for the amino terminal determination of the whole subunit or, for residues identified from the polypeptide cleavage products, above the double-headed arrows spanning the length of the particular polypeptide. One listed pair of polypeptides, V18A and V18B, was sequenced as a mixture and, therefore, are only confirmatory of the cDNA-deduced amino acid sequence, see Figures 1 and 5.

[0021] Samples of the reduced and carboxymethylated pure VEGF AB, A and B monomers, were each digested with the Lys-C endoproteinase, which cleaves polypeptides on the C-terminal side of lysine residues. The peptides were isolated by reverse phase HPLC and their amino acid sequences were determined as described above. The locations of

the peptides in the final VEGF AB, A and B sequences are shown in Figure 2 and Figure 3, respectively.

[0022] The full length coding region of the A subunit or monomer is determined from three sets of overlapping cDNA clones. Degenerate oligonucleotide primers based on the amino acid sequences Phe-Met-Asp-Val-Tyr-Gln from polypeptide L42 (residues 42-47) and Cys-Lys-Asn-Thr-Asp from polypeptide T38 (residues 164-168) (see Figure 1) were used to PCR amplify the central region of the cDNA for VEGF A chain following the procedure of Saiki *et al.*, Science 230: 1350-1354 (1985). A single band migrating at 420 bp was gel purified, digested with Sall, ligated into pGEM3Zf(+) and sequenced. The nucleotide sequence obtained (p4238) was used to design antisense and sense PCR primers to amplify the 5' and 3' ends of the cDNA according to the protocol described by Frohman *et al.* Proc. Natl. Acad. Sci. USA 85:8998-9002 (1988). These 5' and 3' clones are denoted p5-15 and pW3, respectively. Regions of complete DNA sequences, excluding the primers, determined for each set of clones are indicated by double-headed arrows above the nucleotide sequence. In addition to the cDNA coding the 164 amino acid secreted form identified by protein sequencing, two alternatively spliced cDNAs encoding a 146 amino acid and a 214 amino acid forms are cloned and sequenced, Figures 4, 5 and 6.

[0023] The full length coding region of the B subunit or monomer is determined from four sets of overlapping cDNA clones. Degenerate oligonucleotide primers based on the amino acid sequences from polypeptide L50 are used to PCR amplify the central region of the cDNA for VEGF AB, B monomer, following the procedure of Saiki *et al.*, Science 230: 1350-1354 (1985). A single band migrating at 108 bp was gel purified, digested with Sall, ligated into pGEM3Zf(+) and sequenced. The nucleotide sequence obtained (pYG) was used to design antisense and sense PCR primers to amplify the 5' and 3' ends of the cDNA according to the protocol described by Frohman *et al.* Proc. Natl. Acad. Sci. USA 85:8998-9002 (1988). These 5' and 3' clones are denoted p5V2 and p3V2, respectively. Additional 5' end sequences are determined from clone 202 isolated from a cDNA library prepared from GS-9L poly A<sup>+</sup> RNA. Regions of complete DNA sequences, excluding the primers, determined for each set of clones are indicated by double-headed arrows above the nucleotide sequence. The entire base sequence for the 158 amino acid microheterogeneous B subunit and the 138 amino acid microheterogeneous B subunit are shown in Figures 7 and 8.

[0024] The full length coding region of the C subunit or monomer is determined from three sets of overlapping cDNA clones. Degenerate oligonucleotide primers based on the amino acid sequence Phe Ser Pro Ser Cys Val and Glu Met Thr Phe Ser Gly from rat VEGF B subunit are used to PCR amplify the central region of the cDNA of VEGF C chain following the procedure of Saiki *et al.*, Science 230: 1350-1354 (1985). A band migrating at 180 bp is gel purified, reamplified and digested with Sall, ligated into pGEM3Zf(+) and sequenced. The nucleotide sequence obtained (pFSEM') is used to design antisense and sense PCR primers to amplify the 5' and 3' ends of the cDNA according to the protocol described by Frohman *et al.*, Proc. Natl. Acad. Sci. USA 85: 8998-9002 (1988). The 5' and 3' clones are denoted p5:16 and p3:19, respectively. The entire base sequence and amino acid sequence for the C subunit are shown in Figure 9.

[0025] It is intended that vascular endothelial cell growth factor of the present invention exist as a heterodimer consisting of an A microheterogeneous and/or alternatively spliced subunit or a B microheterogeneous and/or alternatively spliced subunit combined with a C microheterogeneous and/or alternatively spliced subunit. It is further intended that VEGF homodimer of the present invention exist as two C subunits. The native forms of the A, B, C subunits may be processed form alternatively spliced full length translation products. The heterodimers or heterodimeric species can be depicted as: A+C or B+C with the A, B or C subunits existing in any of the alternatively spliced or microheterogeneous forms. The homodimers or homodimeric species can be formed by combinations of any of the alternatively spliced or microheterogeneous forms. It is also intended that the invention include all of the individual subunit forms of the C subunit of VEGF.

[0026] It is further intended that the nucleotide sequence for vascular endothelial cell growth factor be interpreted to include all codons that code for the appropriate amino acids in the sequence for each of the vascular endothelial growth factor subunits, as indicated by the degeneracy of the genetic code. It is further intended that the nucleotide sequence and the amino acid sequence for VEGF subunits include truncated genes or proteins that result in proteins which exhibit its biological activity similar to vascular endothelial cell growth factor. The scope of the invention is intended to include all naturally occurring mutations and allelic variants and any randomly generated artificial mutants which may change the sequences but do not alter biological activity as determined by the ability to stimulate the division of vascular endothelial cells.

[0027] The above described heterodimers, homodimers and subunits of vascular endothelial cell growth factor are characterized by being the products of chemical synthetic procedures or of procaryotic or eucaryotic host expression of the DNA sequences as described herein. A monomer is defined as a subunit that is not incorporated in an oligomeric unit. Expression of the recombinant VEGF genes (recombinant DNA) is accomplished by a number of different host cells which contain at least one of a number of expression vectors. Expression vectors are defined herein as DNA sequences that are required for the transcription of cloned copies of recombinant DNA sequences or genes and the translation of their mRNAs in an appropriate host. Such vectors can be used to express genes in a variety of hosts such as bacteria, bluegreen algae, yeast cells, insect cells, plant cells and animal cells, with mammalian cells being preferred. The genes may also be expressed using any of a number of virus expression systems. Specifically designated

vectors allow the shuttling of DNA between bacteria-yeast, bacteria-plant or bacteria-animal cells. An appropriately constructed expression vector should contain: an origin of replication for autonomous replication in host cells, selective markers, a limited number of useful restriction enzyme sites, a high copy number, strong promoters and efficient translational stop signals. A promoter is defined as a DNA sequence that directs RNA polymerase to bind to DNA and to initiate RNA synthesis. A strong promoter is one which causes mRNAs to be initiated at high frequency. Expression vectors may include, but are not limited to, cloning vectors, modified cloning vectors, specifically designed plasmids or viruses and cosmids. The expression of mammalian genes in cultured mammalian cells is well known in the art. Sambrook *et al.*, Molecular Cloning, A Laboratory Manual, 2nd Edition, Book 3, Cold Springs Harbor Laboratory Press (1989) and Current Protocols In Molecular Biology, Ausubel *et al.* Eds, Greene Publishing Associates and Wiley-Interscience, 1987 and supplements, disclose various mammalian expression vectors and vector systems along with methods for the introduction of recombinant vectors into mammalian cells. The cDNA for the monomeric forms of the A, B and C subunits can be expressed in a system such as that described by Linemeyer *et al.*, European Patent Application, Publication No. 259,953. The cDNA is incorporated into a commercially available plasmid such as pKK 223-3 (Pharmacia) as modified as by Linemeyer *et al.* and expressed in *E. coli*. Other expression systems and host cells are well known in the art.

[0028] The high Cys content and glycosylation sites of the A, B and C subunits along with the structure of the homo- and heterodimers suggest that expression of biologically active proteins can be carried out in animal cells. Expression may be carried out in Chinese hamster ovary (CHO) cells with the cloned VEGF DNA cotransfected with the gene encoding dihydrofolate reductase (*dhfr*) into *dhfr*<sup>-</sup> CHO cells, see Sambrook *et al.* Transformants expressing *dhfr* are selected on media lacking nucleosides and are exposed to increasing concentrations of methotrexate. The *dhfr* and VEGF genes are thus coamplified leading to a stable cell line capable of expressing high levels of VEGF. The plasmid is designed to encode either an A subunit, a B subunit or a C subunit or a combination of any two of these subunits. The two cDNAs are operably attached so that the protein produced will be dimeric and will have VEGF biological activity. Operably attached refers to an appropriate sequential arrangement of nucleotide segments, cDNA segments or genes such that the desired protein will be produced by cells containing an expression vector containing the operably attached genes, cDNA segments or nucleotides. Plasmids containing a single subunit species may be used to cotransfect a suitable cell line.

[0029] The expressed proteins (homodimers or heterodimers) are isolated and purified by standard protein purification processes. It is to be understood that the expression vectors capable of expressing heterodimeric forms of VEGF will contain two DNA sequences which will encode either an A subunit and/or a DNA sequence which will encode a B subunit with a DNA sequence which will encode a C subunit. Expression vectors capable of expressing homodimeric forms of VEGF will contain either one or two DNA sequences which encode two C subunits.

[0030] The ability of the various species of VEGF to stimulate the division of vascular endothelial cells makes this protein in all microheterogeneous forms and alternative splicing forms useful as a pharmaceutical agent. The protein as used herein is intended to include all microheterogeneous forms as previously described. The protein can be used to treat wounds of mammals including humans by the administration of the novel protein to patients in need of such treatment.

[0031] The novel method for the stimulation of vascular endothelial cells comprises treating a sample of the desired vascular endothelial cells in a nutrient medium with mammalian VEGF, preferably human or rat, at a concentration of about 1-10 ng/ml. If the vascular endothelial cell growth is conducted *in vitro*, the process requires the presence of a nutrient medium such as DMEM or a modification thereof and a low concentration of calf or bovine serum such as about 0 to 2% by volume. Preservatives such as antibiotics may also be included; these are well known in the art.

[0032] The novel growth factors of this invention are useful for the coverage of artificial blood vessels with vascular endothelial cells. Vascular endothelial cells from the patient would be obtained by removal of a small segment of peripheral blood vessel or capillary-containing tissue and the desired cells would be grown in culture in the presence of VEGF and any other supplemental components that might be required for growth. After growth of adequate numbers of endothelial cells in culture to cover a synthetic polymeric blood vessel the cells would be plated on the inside surface of the vessel, such as fixed umbilical vein, which is then implanted in the patient. Alternatively, tubular supports are coated *in vitro* with VEGF prior to implantation into a patient. Following implantation endothelial cells migrate into and grow on the artificial surface. Prior coating of the artificial vessel either covalently or noncovalently, with proteins such as fibrin, collagen, fibronectin or laminin would be performed to enhance attachment of the cells to the artificial surface. The cell-lined artificial vessel would then be surgically implanted into the patient and, being lined with the patient's own cells, would be immunologically compatible. The non-thrombogenic endothelial cell lining should decrease the incidence of clot formation on the surface of the artificial vessel and thereby decrease the tendency of vessel blockage or embolism elsewhere.

[0033] The novel proteins are also used for the production of artificial vessels. Vascular endothelial cells and smooth muscle cells from the patient would be obtained and grown separately in culture. The endothelial cells would be grown

in the presence of VEGF as outlined above. The smooth muscle would be grown in culture by procedures well known in the art. A tubular mesh matrix of a biocompatible polymer (either a synthetic polymer, with or without a coating of proteins, or a non-immunogenic biopolymeric material such as surgical suture thread) would be used to support the culture growth of the smooth muscle cells on the exterior side and vascular endothelial cells on the interior surface. Once the endothelial cells form a confluent monolayer on the inside surface and multiple layers of smooth muscle cells cover the outside, the vessel is implanted into the patient.

[0034] The novel peptides can also be used for the induction of tissue repair or growth. The pure VEGF would be used to induce and promote growth of tissue by inducing vascular growth and /or repair. The peptide can be used either topically for tissue repair or intravascularly for vascular repair. For applications involving neovascularization and healing of surface wounds the formulation would be applied directly at a rate of about 10 ng to about 1 mg/cm<sup>2</sup>/day. For vascular repair VEGF is given intravenously at a rate of about 1 ng to about 100 µg/kg/day of body weight. For internal vascular growth, the formulation would be released directly into the region to be neovascularized either from implanted slow release polymeric material or from slow release pumps or repeated injections. The release rate in either case is about 10 ng to about 100 µg/day/cm<sup>3</sup>.

[0035] For non-topical application the VEGF is administered in combination with pharmaceutically acceptable carriers or diluents such as, phosphate buffer, saline, phosphate buffered saline, Ringer's solution, and the like, in a pharmaceutical composition, according to standard pharmaceutical practice. For topical application, various pharmaceutical formulations are useful for the administration of the active compound of this invention. Such formulations include, but are not limited to, the following: ointments such as hydrophilic petrolatum or polyethylene glycol ointment; pastes which may contain poms such as xanthan gum; solutions such as alcoholic or aqueous solutions; gels such as aluminum hydroxide or sodium alginate gels; albumins such as human or animal albumins; collagens such as human or animal collagens; celluloses such as alkyl celluloses, hydroxy alkyl celluloses and alkylhydroxyalkyl celluloses, for example methylcellulose, hydroxyethyl cellulose, carboxymethyl cellulose, hydroxypropyl methylcellulose, and hydroxypropyl cellulose; polyoxamers such as Pluronic® Polyols exemplified by Pluronic® F-127; tetronics such as tetronic 1508; and alginates such as sodium alginate.

[0036] The following examples illustrate the present invention without, however, limiting the same thereto.

#### EXAMPLE 1

##### Preparation of Medium Conditioned By GS-9L Cells

[0037] GS-9L cells were grown to confluence in 175 cm<sup>2</sup> tissue culture flasks in Dulbecco's Modified Eagle's Medium/10% newborn calf serum (DMEM/NCS). At confluence the medium was decanted from the flasks, the flasks were washed with calcium and magnesium free phosphate buffered saline (PBS) and the cells were removed by treatment with a 1X solution of trypsin/EDTA (Gibco). The cells ( $1 \times 10^8$ ) were pelleted by centrifugation, resuspended in 1500 ml of DMEM/5% NCS and plated into a ten level (6000 cm<sup>2</sup> surface area) cell factory (NUNC). After 72 hours incubation at 37° C in a 5% CO<sub>2</sub> atmosphere the medium was decanted and the cell factories were washed 3 times with PBS. The cells were refed with 1500 ml of a 1:2 mixture of Ham's F-12/DMEM containing 25 mM Hepes, pH 7.4, 5 µg/ml insulin, 10 µg/ml transferrin and 1.0 mg/ml bovine serum albumin. This medium was changed with fresh F-12/DMEM after 24 hours and collected every 48 hours after that. The conditioned medium was filtered through a Whatman #1 paper to remove cell debris and stored frozen at -20°C.

#### EXAMPLE 2

##### Carboxymethyl-Sephadex Chromatography of VEGF AA and VEGF AB

[0038] GS-9L conditioned medium, from Example 1, was thawed and brought to pH 6.0 with 1 M HCl. Two grams of CM Sephadex C-50 cation exchange (Pharmacia) resin preequilibrated in PBS adjusted to pH 6.0 with 1 N HCl were added to 20 liters of conditioned medium. The mixture was stirred at low speed for 24 hours at 4° C. The resin was then allowed to settle and the medium was siphoned off. The remaining resin slurry was packed into a 3.0 cm diameter column and any remaining medium was allowed to drain off. Unbound protein was washed off the column with 0.05 M sodium phosphate, pH 6.0, containing 0.15 M NaCl. Vascular endothelial growth factor activity was eluted from the column with a subsequent wash of 0.05 M sodium phosphate, pH 6.0, containing 0.6 M NaCl.



## EXAMPLE 3

Concanavalin A (Con A) Lectin Affinity Chromatography of VEGF AA and VEGF AB

5 [0039] A 0.9 cm diameter column containing about 5 ml of packed Con A agarose (Vector Laboratories) was equilibrated with 0.05 M sodium acetate, pH 6.0, containing 1 mM  $\text{Ca}^{++}$ , 1 mM  $\text{Mn}^{++}$  and 0.6 M NaCl. The active eluate from the CM Sephadex C-50 column, Example 2, was applied to the Con A agarose and unbound protein was washed from the column with equilibration buffer. The column was then rinsed with three column volumes of 0.05 M sodium acetate, pH 6.0, containing 1 mM  $\text{Ca}^{++}$ , 1 mM  $\text{Mn}^{++}$  and 0.1 M NaCl. Bound protein was subsequently eluted from the column  
 10 by application of this buffer supplemented with 0.32 M  $\alpha$ -methyl mannoside and 0.28 M  $\alpha$ -methyl glucoside.

## EXAMPLE 4

Polyaspartic Acid WCX HPLC Cation Exchange Chromatography of VEGF AA and VEGF AB

15 [0040] The active eluate from the Con A column, Example 3, was applied to a 25 cm x 4.6 mm poly(aspartic acid) WCX cation exchange HPLC column (Nest Group) pre-equilibrated in 0.05 M sodium phosphate buffer, pH 6.0. The column was eluted with a linear gradient of 0 to 0.75 M NaCl in this buffer over 60 minutes at a flow rate of 0.75 ml/min collecting 0.75 ml fractions. VEGF AB activity present in fractions eluting between approximately 21.7 and 28.5 ml were  
 20 pooled.

## EXAMPLE 5

Metal Chelate Chromatography

25 [0041] The active fractions eluted from the poly(aspartic acid) WCX column, Example 4, that contain VEGF AB were pooled, adjusted to pH 7.0 and loaded onto a 1 x 10 cm column of Pharmacia Chelating Sepharose 6B charged with an excess of copper chloride and equilibrated in 0.05 M sodium phosphate, pH 7.0, containing 2 M NaCl and 0.5 mM imidazole (A buffer). VEGF AB was eluted from the column with a gradient from 0-20% B over 10 minutes, 20-35% B  
 30 over 45 minutes and 35-100% B over 5 minutes at a flow rate of 0.3 ml/min, where B buffer was 0.05 M sodium phosphate, pH 7.0, containing 2 M NaCl and 100 mM imidazole. The active fractions containing VEGF AB activity eluting between 12.6 and 22.8 ml of the gradient effluent volume were pooled.

## EXAMPLE 6

Reverse Phase Chromatography

35 [0042] The fractions containing VEGF AB activity pooled from the metal chelate column, Example 5 were loaded onto a 4.6 mm x 5 cm Vydac  $\text{C}_4$  reverse phase HPLC column (5  $\mu\text{m}$  particle size) equilibrated in solvent A (0.1% trifluoroacetic acid (TFA)). The column was eluted with a gradient of 0-30% solvent B over 15 minutes, 30% B for an additional  
 40 15 minutes, then 30-45% B over 22.5 minutes and finally 45-100% B over 5.5 minutes where solvent B = A containing 67% acetonitrile. The flow rate was maintained at 0.75 ml/min. The active VEGF AB fractions eluting between approximately 32.2 and 37.5 ml of the gradient effluent volume were pooled.

## 45 EXAMPLE 7

Mitogenic Assays

50 [0043] Human umbilical vein endothelial cells (HUVE) were plated on gelatin-coated 48 well tissue culture dishes at a density of 5000 cells/well in 500  $\mu\text{l}$  of Medium 199 containing 20% heat inactivated fetal calf serum (FCS). Samples to be assayed were added at the time of plating. The tissue culture plates are incubated at 37° C for 12 hours and 2 microcuries of tritiated thymidine (NEN, 20 Ci/mmol) was added per ml of assay medium (1.0  $\mu\text{Ci}/\text{well}$ ). The plates were incubated for a further 60 hr, the assay medium was removed and the plates were washed with Hanks balanced salt solution containing 20 mM Hepes, pH 7.5, and 0.5 mg/ml bovine serum albumin. The cells were lysed and the labelled  
 55 DNA solubilized with 200  $\mu\text{l}$  of a solution containing 2 gm of sodium carbonate and 400 mg sodium hydroxide in 100 ml water. The incorporated radioactivity was determined by liquid scintillation counting.

[0044] The concentration of VEGF AB which elicited a half-maximal mitogenic response in HUVE cells was approximately  $2 \pm 1$  ng/ml. The glycosaminoglycan heparin, which is required in these assays at a level of 10-100  $\mu\text{g}/\text{ml}$  to pro-

mote a response to a positive control, acidic fibroblast growth factor, does not enhance mitogenic stimulation of these cells by VEGF AB.

#### EXAMPLE 8

##### Purity And Protein Structural Characterization of VEGF AB

[0045] Purity of the protein under non-reducing conditions was determined by SDS-PAGE in 12.5% crosslinked gels according to the method of Laemmli, Nature 227: 680-685 (1970). The silver-stained gel contained a single band with an apparent mass of approximately 58 kDa. VEGF AB migrated in SDS-PAGE under reducing conditions in 15% crosslinked gels as a broad silver-stained band with apparent molecular mass of approximately 23 kDa.

[0046] VEGF AB was stored at 4°C in the aqueous trifluoroacetic acid (TFA)/acetonitrile mixture used to elute the homogeneous protein in reversed phase C<sub>4</sub> HPLC chromatography at the final stage of the purification protocol previously described. Aliquots of the purified protein (1-2 µg) were vacuum evaporated to dryness in acid-washed 10 x 75 mm glass tubes and reduced for 2 hours at 50°C in 100 µl of 0.1 M Tris buffer, pH 9.5, and 6 M guanidinium chloride containing 0.1% EDTA and 20 mM dithiothreitol (Calbiochem, Ultrol grade) under an argon atmosphere. The reduced protein was subsequently carboxymethylated for 1 hour at 20°C by the addition of 100 µl of 0.7 M Tris, pH 7.8, containing 0.1% EDTA, 6 M guanidinium chloride, 9.2 µM unlabeled iodoacetic acid and 50 µCi of iodo[2-<sup>14</sup>C]acetic acid (17.9 mCi/mole, Amersham). After completion of the carboxymethylation, the mixture was loaded directly onto a 4.6 mm x 5.0 cm Vydac C<sub>4</sub> column which had been preequilibrated in 0.1% TFA. The reduced and carboxymethylated protein was repurified by elution with a 45 minute linear gradient of 0 to 67% (v/v) acetonitrile in 0.1% TFA at a flow rate of 0.75 ml/min and stored in this elution solution at 4°C. The reduced and carboxymethylated protein eluted as two peaks at approximately 23 and 25 ml that were of approximately equal area as determined by monitoring absorbance at 210 nm.

[0047] Samples of the two protein subunits isolated after reduction and carboxymethylation were each applied to polybrene-coated glass fiber filters and their N-terminal sequences were determined by Edman degradation in an ABI gas phase microsequencer in conjunction with an ABI 120A on line phenylthiohydantoin analyzer following manufacturers instructions. The peak of absorbance eluting at approximately 25 ml (A subunit) yielded an amino terminal sequence Ala Pro Thr Thr Glu Gly Glu Gln Lys Ala His Glu Val Val SEQ ID NO:1 identical to VEGF AA. The peak of absorbance eluting at approximately 23 ml (B subunit) yielded the N-terminal sequence Ala Leu Ser Ala Gly Asn Xaa Ser Thr Glu Met Glu Val Val Pro Phe Asn Glu Val SEQ ID NO:2 plus a nearly equal amount of a truncated form of the same sequence missing the first three residues. The missing X residue corresponds to an Asn in the cloned sequence. Since this missing Asn occurs in a classical Asn-X-Ser/Thr N-glycosylation sequence it is presumed to be glycosylated. The A and sum of the B chain peptides were recovered in nearly equal amounts supporting the interpretation that the two peptides combine to form an AB heterodimer in VEGF II.

[0048] Reduced and carboxymethylated A and B subunits (650 ng each) were each dried by vacuum evaporation in acid-washed 10 x 75 mm glass tubes. Lys C protease (50 ng, Boehringer Mannheim), an enzyme that cleaves on the carboxyl terminal side of lysine residues, was added to each tube in 100 µl of 25 mM Tris, pH 8.5, 0.1% EDTA. The substrate protein subunits were separately digested at 37°C for 8 hours and the resulting polypeptides resolved by reversed phase HPLC chromatography on a 4.6 mm x 25 cm Vydac C<sub>18</sub> column equilibrated in 0.1% TFA. Polypeptides were fractionated by elution with a 2 hour linear gradient of 0-67% acetonitrile in 0.1% TFA at a flow rate of 0.75 ml/min at 20°C. Individual peaks were manually collected and stored in this elution solution at 4°C.

[0049] The amino acid sequences of the isolated peptides were then determined using Edman degradation in an ABI gas phase sequencer in conjunction with the ABI 120 A on line phenylthiohydantoin analyzer (Applied Biosystems Int.). The peptide sequences are shown in the following Figures 2 and 3. The amino acid sequence of Lys C fragment L20 (Fig. 5) demonstrates that the form of VEGF AB mature A subunit in the heterodimer is the 164 amino acid form. The amino acid sequence of Lys C fragment L26 (Fig. 3) demonstrates that the form of VEGF AB mature B subunit in the heterodimer is the 135 amino acid form derived from the 158 full length amino acid form.

#### EXAMPLE 9

##### Cloning and Sequencing of the VEGF A Monomer

##### PCR Amplification, Cloning and Sequencing of P4238

[0050] Two degenerate oligonucleotides were synthesized in order to amplify the cDNA encoding the peptide sequences of VEGF A subunit between LysC fragment L 42 and tryptic fragment T38. These oligonucleotides were:

**L42.2 5' TTTGTCGACTT[TC]ATGGA[TC]GT[N]TA[TC]CA 3'**  
**SEQ ID NO:3**

5

**T383'B**

**5' CAGAGAATTTCGTCGACA[AG]TC[N]GT[AG]TT[TC]TT**  
**[AG]CA 3' SEQ ID NO:4**

10

where N=ACGT

[0051] Poly A<sup>+</sup> RNA was isolated from GS-9L cells using the Fast Track RNA isolation kit from Invitrogen and the protocol provided. First strand cDNA synthesis was performed as follows;

15 1 µg of GS-9L RNA was annealed to 1 µg of adapter primer TA17,  
 5' GACTCGAGTCGACATCGATTTTTTTTTTTTTTTTTT 3' SEQ ID NO:5, by incubating in a volume of 10 µl at 70°C for 5 min. followed by cooling to room temperature. To this reaction was added:

20	3.0 µl	water
	2.5 µl	10X buffer (500 mM Tris-HCl pH 8.3, 750 mM KCl, 100mM MgCl <sub>2</sub> , 5 mM spermidine)
	2.5 µl	100 mM DTT
	2.5 µl	10 mM each dATP, dGTP, dCTP, dTTP
	0.6 µl	15 units RNasin
	2.5 µl	40 mM Na pyrophosphate
25	1.5 µl	15 units reverse transcriptase

and the reaction was incubated at 42°C for 1 hour, then diluted to 1 ml in 10 mM Tris-HCl 1 mM EDTA, pH 7.5.

PCR Reactions:

30

Primary reaction (100 µl)

[0052]

35	10 µl	10X buffer from Perkin Elmer Cetus GeneAmp kit
	16 µl	1.25 mM each stock of dATP, dCTP, dGTP, and dTTP
	2 µl	first strand GS9L cDNA
	2 µl	50 pMoles L42.2
	2 µl	50 pMoles T383'B
40	0.5 µl	2.5 units Amplitaq DNA polymerase
	67.5 µl	water

[0053] Reaction conditions, 40 cycles of 94°C, 1'; 50°C, 2'30"; 72°C, 2'.

45 Prep scale secondary reaction:

[0054]

	100 µl	10X buffer
50	160 µl	1.25 mM each stock of dATP, dCTP, dGTP, and dTTP
	10 µl	primary PCR reaction
	20 µl	500 pMoles L42.2
	20 µl	500 pMoles T383'B
	5 µl	25 units Amplitaq DNA polymerase
55	685 µl	water

[0055] Reaction conditions 94°C, 1'; 55°C, 2'; 72°C, 2'; 30 cycles.

[0056] The PCR product was concentrated by Centricon 30 spin columns, purified on a 1% agarose gel, and digested

with restriction endonuclease Sall. The Sall fragment was then ligated into Sall cut pGEM3Zf(+). The ligation mix was used to transform *E. coli* XL-1 blue. Plasmid DNA was isolated from white transformants and sequenced by the dideoxy chain termination method.

#### 5 PCR Amplification, Cloning and Sequencing of pW-3

[0057] Based on the sequence obtained from the p4238 clones, two specific PCR primers were synthesized; oligo 307 5' TTTGTCGACTCAGAGCGGAGAAAGC 3' SEQ ID NO:6 and oligo 289 5' TTTGTCGACGAAAATCACTGTGAGC 3' SEQ ID NO:7. These primers were used in combination with oligoA 17

- 10 5' GACTCGAGTCGACATCG 3' SEQ ID NO:8 to amplify the cDNA encoding the COOH terminus of VEGF A subunit using the 3' RACE technique described by Frohman *et al.*, PNAS 85: 8998-9002 (1988).

PCR reactions:

- 15 Primary reaction 100  $\mu$ l

[0058]

- 10  $\mu$ l 10X buffer from Perkin Elmer Cetus GeneAmp kit  
20 18  $\mu$ l 1.25 mM each stock of dATP, dCTP, dGTP, and dTTP  
0.35  $\mu$ l first strand GS-9L cDNA  
2  $\mu$ l 50 pMoles oligo 289  
0.5  $\mu$ l 2.5 units Amplitaq DNA polymerase  
67.15  $\mu$ l water

25 [0059] Reaction conditions 94°C, 1'; 58°C, 2'; 72°C, 2'; 10 cycles then add 50 pMoles A17, then 1 cycle of 94°C, 1'; 58°C, 2'; 72°C, 40' followed by 40 cycles 94°C, 1'; 58°C, 2'; 72°C, 2'.

Prep Scale secondary reaction:

30 [0060]

- 60  $\mu$ l 10X buffer  
108  $\mu$ l 1.25 mM each stock of dATP, dCTP, dGTP, and dTTP  
35 24  $\mu$ l primary PCR reaction  
12  $\mu$ l 300 pMoles oligo 307  
12  $\mu$ l 300 pMoles oligo A17  
3  $\mu$ l 15 units Amplitaq DNA polymerase  
381  $\mu$ l water

40 [0061] Reaction conditions 94°C, 1'; 58°C, 2'; 72°C, 2'; 30 cycles.

[0062] The PCR product was purified on a 1% agarose gel and digested with restriction endonuclease Sall. The Sall fragment was then ligated into Sall cut pGEM3Zf(+). The ligation mix was used to transform *E. coli* XL-1 blue. Plasmid DNA was isolated from white transformants and sequenced by the dideoxy chain termination method.

#### 45 PCR Amplification, Cloning and Sequencing of p5-15

- [0063] Based on the sequence of p4238 clones, two specific PCR primers were synthesized; oligo 113 5' TTTGTCGACAAACAGGACGGCTTGAAG 3' SEQ ID NO:9 and oligo 74 5' TTTGTCGACATACTCCTGGAAGATGTCC 3' SEQ ID NO:10. These primers were used in combination with oligo A17 5' GACTCGAGTCGACATCG 3' SEQ ID NO:8 to amplify the cDNA encoding the amino terminus of VEGF A subunit using the 5' RACE technique described by Frohman *et al.*, supra. Oligo 151 was synthesized in order to specifically prime VEGF A subunit cDNA from GS-9L RNA. Oligo 151 is 5' CTTCATCATTGCAGCAGC 3' SEQ ID NO:11.

55 [0064] RNA was isolated from GS-9L cells using the Fast Track RNA isolation kit from Invitrogen using the protocol provided. First strand cDNA synthesis was performed as follows;

[0065] One  $\mu$ g of GS9L RNA was annealed to 1  $\mu$ g of oligo 151 by incubating in a volume of 6  $\mu$ l at 70°C for 5' followed by cooling to room temperature. To this reaction was added:

1.5 $\mu$ l	10X buffer (500mM Tris-HCl, pH 8.3, 750 mM KCl, 100 mM MgCl <sub>2</sub> , 5 mM spermidine)
2.5 $\mu$ l	10 mM DTT
2.5 $\mu$ l	10 mM each dATP, dGTP, dCTP, dTTP
0.6 $\mu$ l	25 units RNasin
5 2.5 $\mu$ l	40 mM Na pyrophosphate
9.5 $\mu$ l	20 units diluted reverse transcriptase

[0066] The reaction was incubated at 42°C for 1 hour. Excess oligo151 was removed by Centricon 100 spin columns and the 5' end of the cDNA was tailed by the addition of dATP and terminal transferase. The tailed cDNA was diluted to  
10 a final volume of 150  $\mu$ l in 10 mM Tris-HCl, 1 mM EDTA, pH 7.5.

#### PCR Reactions:

##### Primary reaction (50 $\mu$ l)

15

##### [0067]

5 $\mu$ l	10X buffer from Perkin Elmer Cetus GeneAmp Kit
8 $\mu$ l	1.25 mM each stock of dATP, dCTP, dGTP, and dTTP
20 5 $\mu$ l	first strand GS-9L cDNA prime with oligo 151 and tailed
1 $\mu$ l	25 pMoles oligo 113
1 $\mu$ l	25 pMoles oligo A17
1 $\mu$ l	10 pMoles oligo TA17
0.25 $\mu$ l	1.25 units Ampliqa DNA polymerase
25 28.75 $\mu$ l	water

[0068] Reaction conditions; 1 cycle 94°C 1'; 50°C 2'; 72°C 40' then 40 cycles of 94°C 1'; 50°C 1'30"; 72°C 2'

##### Prep scale secondary reaction:

30

##### [0069]

60 $\mu$ l	10X buffer
96 $\mu$ l	1.25 mM each stock of dATP, dCTP, dGTP, and dTTP
35 6 $\mu$ l	primary PCR reaction
12 $\mu$ l	300 pMoles oligo74
12 $\mu$ l	300 pMoles oligo A17
3 $\mu$ l	15 units Ampliqa DNA polymerase
411 $\mu$ l	water

40

[0070] Reaction conditions 94°C, 1'; 55°C, 2'; 72°C, 2' 30 cycles.

[0071] The PCR product was concentrated by Centricon 100 spin columns, and digested with restriction endonuclease Sall. The Sall fragment was then ligated into Sall cut pGEM3Zf(+). The ligation mix was used to transform E. coli XL-1 blue. Plasmid DNA was isolated from white transformants and sequenced by the dideoxy chain termination  
45 method. The base sequence is shown in Fig. 5.

#### Cloning and sequencing of alternative forms of VEGF A cDNA

[0072] Based on the sequence obtained from the p5-15 and pW-3 clones, two specific PCR primers were synthesized; oligo 5'C 5' TTTGTCGACAACCATGAACTTTCTGC 3' SEQ ID NO:12 and oligo 181 5' TTTGTCGACGGTGA-GAGGTCTAGTTC 3' SEQ ID NO:13. These primers were used together to amplify multiple cDNAs encoding alternative forms of the VEGF A subunit.

55

##### Preparative PCR Reaction:

##### [0073]

50 $\mu$ l	10X buffer
------------	------------

80 µl	1.25mM each stock of dATP, dCTP, dGTP, and dTTP
10 µl	first strand GS-9L cDNA
10 µl	300pMoles oligo 5'C
10 µl	300pMoles oligo 181
5 2.5 µl	15 units Amplitaq DNA polymerase
337.5 µl	water

[0074] Reaction conditions 94°C, 1'; 58°C, 2'; 72°C, 3'; 40 cycles.

[0075] The PCR product was extracted with phenol/chloroform, concentrated by Centricon 30 spin columns, precipitated by ethanol, and digested with restriction endonuclease Sal I, and ligated into SalI cut pGEM3Zf(+). The ligation mix was used to transform *E. coli* XL-1 blue. Plasmid DNA was isolated from white transformants and sequenced by the dideoxy chain termination method. Three sets of clones were identified. Clone#12 encoded the 190 amino acid form of VEGF A subunit identical to that shown in Fig. 1. The 164 amino acid secreted form of VEGF A subunit is that amino acid sequence running continuously from Ala<sup>27</sup> to Arg<sup>190</sup>. Clone#14 has a 135 base pair deletion between the second base of the Asn<sup>140</sup> codon and the third base of the Arg<sup>184</sup> codon. This clone thus encodes a 146 aa form of the VEGF A subunit with the conversion of Asn<sup>140</sup> to Lys<sup>140</sup>. The 120 amino acid secreted form of VEGF A subunit runs from Ala<sup>27</sup> to Asn<sup>140</sup>, which becomes Lys<sup>140</sup> and does not begin until Cys<sup>185</sup>, this form also finishes at Arg<sup>190</sup>, Figure 4. Clone #16 has a 72 base pair insertion between the second and third base of the Asn<sup>140</sup> codon. This clone thus encodes the 214 amino acid form of the VEGF A subunit with the conversion of Asn<sup>140</sup> to Lys<sup>140</sup>, Figure 6.

#### EXAMPLE 10

##### Cloning and Sequencing of the VEGF B Subunit

##### 25 PCR Amplification, Cloning and Sequencing of pYG

[0076] Two degenerate oligonucleotides were synthesized in order to amplify the cDNA encoding the peptide sequences of VEGF B on Lys C fragment L50. These oligonucleotides were:

YI 5' TTTGTCGACATA[TC]AT[TC]GC[N]GA[TC]GA[AG]C 3' SEQ ID NO:14

30 GC 5' TTTGTCGACTC[AG]TC[AG]TT[AG]CA[AG]CA[N]CC 3' SEQ ID NO:15 where N=ACGT

[0077] RNA was isolated from GS-9L cells using the Fast Track RNA isolation kit from Invitrogen and the protocol provided. First strand cDNA synthesis was performed as follows;

35 1 µg of GS-9L poly A<sup>+</sup>RNA was annealed to 1 µg of adapter primer TA17,  
5'GACTCGAGTCGACATCGATTTTTTTTTTTTTTTT 3' SEQ ID NO: 5, by incubating in a volume of 10 µl at 70°C for 5 min. followed by cooling to room temperature. To this reaction was added:

3.0 µl	water
2.5 µl	10X buffer (500 mM Tris-HCl, pH 8.3, 750 mM KCl, 100 mM MgCl <sub>2</sub> , 5mM spermidine)
40 2.5 µl	100 mM DTT
2.5 µl	10 mM each dATP, dGTP, dCTP, dTTP
0.6 µl	15 units RNasin
2.5 µl	40 mM Na pyrophosphate
1.5 µl	15 units reverse transcriptase

45 and the reaction was incubated at 42°C for 1 hour, then diluted to 1 ml in 10 mM Tris-HCl, 1 mM EDTA, pH 7.5.

##### PCR Reactions:

##### 50 Primary reaction (50µl)

##### [0078]

5 µl	10X buffer from Perkin Elmer Cetus GeneAmp kit
55 8 µl	1.25 mM each stock of dATP, dCTP, dGTP, and dTTP
1 µl	first strand GS-9L cDNA
1 µl	50 pMoles oligo YI
1 µl	50 pMoles oligo GC

0.25  $\mu$ l 1.25 units Amplitaq DNA polymerase  
33.75  $\mu$ l water

[0079] Reaction conditions, 40 cycles of 94°C, 1'; 50°C, 2'; 72°C, 2'.

Prep scale reaction:

[0080]

60  $\mu$ l 10X buffer  
96  $\mu$ l 1.25mM each stock of dATP, dCTP, dGTP, and dTTP  
12  $\mu$ l first strand 659L cDNA  
12  $\mu$ l 500pMoles oligo YI  
12  $\mu$ l 500pMoles oligo GC  
3  $\mu$ l 15 units Amplitaq DNA polymerase  
405  $\mu$ l water

[0081] Reaction conditions 94°C, 1'; 50°C, 2'; 72°C, 2' 40 cycles.

[0082] The PCR product was concentrated by Centricon 30 spin columns and digested with restriction endonuclease Sall. The Sall fragment was then ligated into Sall cut pGEM3Zf(+). The ligation mix was used to transform *E. coli* XL-1 blue. Plasmid DNA was isolated from white transformants and sequenced by the dideoxy chain termination method.

#### PCR Amplification, Cloning and Sequencing of p3V2

[0083] Based on the sequence obtained from the pYG clones, a specific PCR primer was synthesized; oligo HP 5' TTTGTCGACACACCTAATGAAGTGTC 3' SEQ ID NO:16. This primer was used in combination with oligo A17 5' GACTCGAGTCGACATCG 3' SEQ ID NO:8 to amplify the cDNA encoding the COOH terminus of the VEGF B subunit using the 3' RACE technique described by Frohman *et al.*, PNAS 85: 8998-9002 (1988).

Preparative PCR reaction:

[0084]

60  $\mu$ l 10X buffer from Perkin Elmer Cetus Gene Amp Kit  
12  $\mu$ l first strand 659L cDNA  
96  $\mu$ l 1.25 mM each of dATP, dCTP, dGTP, dTTP  
12  $\mu$ l 300 pMoles oligo A17  
12  $\mu$ l 300 pMoles oligo HP  
3  $\mu$ l 15units Amplitaq DNA polymerase  
405  $\mu$ l water

[0085] Reaction conditions 1 cycle of 94°C, 1'; 58°C, 2'; 72°C, 2'; followed by 40 cycles 94°C, 1', 58°C, 2' and 72°C, 2'.

[0086] The PCR product was concentrated by Centricon 30 spin columns, precipitated with ethanol and digested with restriction endonuclease Sa1I. The Sa1I fragment was then ligated into Sa1I cut pGEM3Zf(+). The ligation mix was used to transform *E. coli* XL-1 blue. Plasmid DNA was isolated from white transformants and sequenced by the dideoxy chain termination method.

#### PCR Amplification, Cloning and Sequencing of p5V2

[0087] Based on the sequence of pYG clones, two specific PCR primers were synthesized; oligoVL' 5' TTTGTCGACACAGCGACTCAGAAGG 3' SEQ ID NO: 17 and oligoVS' 5' TTTGTCGACACTGAATATGAGACAC 3' SEQ ID NO:18. These primers were used in combination with oligo A17 5' GACTCGAGTCGACATCG 3' SEQ ID NO:8 to amplify the cDNA encoding the amino terminus of the VEGF B subunit using the 5' RACE technique described by Frohman *et al.*, supra. Oligo 151 was synthesized in order to prime cDNA from GS-9L RNA. Oligo 151 is 5' CTTTCATTCATTGCAGCAGC 3' SEQ ID NO:11.

[0088] Poly A<sup>+</sup> RNA was isolated from GS9L cells using the Fast Track RNA isolation kit from Invitrogen using the protocol provided. First strand cDNA synthesis was performed as follows:

[0089] One  $\mu$ g of GS9L RNA was annealed to 1  $\mu$ g of oligo 151 by

incubating in a volume of 6  $\mu$ l at 70°C for 5' followed by cooling to room temperature. To this reaction was added:

1.5  $\mu$ l 10X buffer (500 mM Tris-HCl, pH 8.3, 750 mM KCl, 100 mM MgCl<sub>2</sub>, 5mM spermidine)  
 5 2.5  $\mu$ l 10 mM DTT  
 2.5  $\mu$ l 10 mM each dATP, dGTP, dCTP, dTTP  
 0.6  $\mu$ l 25 units RNasin  
 2.5  $\mu$ l 40 mM Na pyrophosphate  
 9.5  $\mu$ l 20 units diluted reverse transcriptase

10 [0090] The reaction was incubated at 42°C for 1 hour.

[0091] Excess oligo 151 was removed by Centricon 100 spin columns and the 5' end of the cDNA was tailed by the addition of dATP and terminal transferase. The tailed cDNA was diluted to a final volume of 150  $\mu$ l in 10 mM Tris-HCl, 1 mM EDTA, pH 7.5

15 PCR Reactions:

Primary reaction (50  $\mu$ l)

20 [0092]

5  $\mu$ l 10X buffer from Perkin Elmer Cetus GeneAmp Kit  
 8  $\mu$ l 1.25 mM each stock of dATP, dCTP, dGTP, and dTTP  
 5  $\mu$ l first strand GS9L cDNA primed with oligo151 and tailed  
 25 1  $\mu$ l 25 pMoles oligo VL'  
 1  $\mu$ l 25 pMoles oligo A17  
 1  $\mu$ l 10 pMoles oligo TA17  
 0.25  $\mu$ l 1.25 units AmpliQ DNA polymerase  
 28.75  $\mu$ l water

30 [0093] Reaction conditions; 1 cycle 94°C, 1'; 58°C, 2'; 72°C, 40' then 40 cycles of 94°C, 1'; 58°C, 2'; 72°C, 2'.

Prep scale secondary reaction:

35 [0094]

100  $\mu$ l 10X buffer  
 160  $\mu$ l 1.25 mM each stock of dATP, dCTP, dGTP, and dTTP  
 10  $\mu$ l primary PCR reaction  
 40 20  $\mu$ l 500 pMoles oligo VS'  
 20  $\mu$ l 300 pMoles oligo A17  
 5  $\mu$ l 25 units AmpliQ DNA polymerase  
 685  $\mu$ l water

45 [0095] Reaction conditions 94°C, 1'; 58°C, 2'; 72°C, 2' 30 cycles.

[0096] The PCR product was extracted with phenol/chloroform, concentrated by Centricon 30 spin columns, precipitated by ethanol, and digested with restriction endonuclease Sall. The Sall fragment was purified on 4% Nu-Sieve Agarose gel then ligated into Sall cut pGEM3Zf(+). The ligation mix was used to transform *E. coli* XL-1 blue. Plasmid DNA was isolated from white transformants and sequenced by the dideoxy chain termination method.

50 PCR Amplification, Cloning and Sequencing of pCV2 and pCV2.1

[0097] Based on the sequences of the p3V2 and p5CV2 clones, two specific PCR primers were synthesized; oligo 5'CV2.1 5' TTTGTCTGAC[N][N]GCAGGTCCTAGCTG 3' SEQ ID NO:19 and oligo 3'CV2 5' TTTGTC-  
 55 GAC[N][N]CTAATAAATAGAGGG 3' SEQ ID NO:20. These primers were used together to amplify the cDNA encoding the VEGF B subunit.



## Preparative PCR Reaction:

[0098]

- 5    40  $\mu$ l    10X buffer  
      64  $\mu$ l    1.25 mM each dATP, dTTP, dGTP, dCTP  
      8  $\mu$ l    first strand GS-9L cDNA  
      8  $\mu$ l    200 pMoles 5'CV2.1  
      8  $\mu$ l    200 pMoles 3'CV2  
 10   2  $\mu$ l    10units Amplitaq DNA polymerase  
      270  $\mu$ l    water

[0099] Reaction conditions: 94°C, 1', 58°C, 2', 72°C, 2'; 40 cycles.

- [0100] The PCR product was extracted with phenol/chloroform, concentrated by Centricon 30 spin columns, precipitated by ethanol, and digested with restriction endonuclease Sal I, and ligated into Sal I cut pGEM3Z(+). The ligation mix was used to transform *E. coli* XL-1 blue. Plasmid DNA was isolated from white transformants and sequenced by the dideoxy chain termination method. Two sets of clones were identified, one encoded a 158 amino acid sequence and the other encoded a 138 amino acid sequence, see Figures 7 and 8.

20    cDNA Cloning of VEGF B Subunit

[0101] The DNA and protein sequences for the amino terminus of the signal peptide of VEGF B was determined from a cDNA clone isolated from a cDNA library constructed from GS-9L polyA<sup>+</sup> RNA.

## 25    First Strand Synthesis

[0102] Anneal 15.6 $\mu$ l (5ug) GS-9L polyA<sup>+</sup> RNA and 2.5 $\mu$ l (2.5ug) oligo dT-XbaI primer by heating to 70° C 5' slow cool to room temperature. Add the following:

- 30   5.5 $\mu$ l    10X buffer (500 mM Tris-HCl, pH 8.3 (42° C), 750 mM KCl, 100 mM MgCl<sub>2</sub>, 5mM spermidine  
      5.5 $\mu$ l    100mM DTT  
      5.5 $\mu$ l    10 mM each dATP, dTTP, dCTP, dGTP  
      1.4 $\mu$ l    (55units) RNasin  
      5.5 $\mu$ l    40mM NaPPi  
 35   13.5 $\mu$ l    55units AMV reverse transcriptase

[0103] Incubate at 42° C 60'.

## Second Strand Synthesis:

40

[0104]

Assemble reaction mix

45    [0105]

- 50  $\mu$ l    first strand reaction  
      25  $\mu$ l    10X buffer ( 500 mM Tris-HCl, pH7.2, 850 mM KCL, 30 mM MgCl<sub>2</sub> 1mg/ml BSA, 100 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>  
      7.5  $\mu$ l    100 mM DTT  
 50   25  $\mu$ l    1mM NAD  
      6.5  $\mu$ l    (65units) *E. coli* DNA Polymerase I  
      2.5  $\mu$ l    (2.5units) *E. coli* DNA Ligase  
      2.5  $\mu$ l    (2 units) *E. coli* RNase H  
      135  $\mu$ l    water

55

[0106] Incubate at 14° C for 2h and then incubate 70° C for 10'. Add 1ul (10 units) T4 DNA Polymerase, incubate at 37° C for 10', add 25  $\mu$ l 0.2M EDTA an extract with phenol/chloroform, then precipitate by the addition of 0.5 volume of 7.5 M ammonium acetate and 3 volumes of ethanol, collect precipitate and resuspend in 20  $\mu$ l of 10 mM Tris-HCl, pH

7.5, 1mM EDTA.

#### cDNA Library Construction

- 5 [0107] The above cDNA was ligated into EcoR1/ XbaI digested LambdaGEM-4 (Promega Biochemicals) after the addition of EcoR1 linkers and digestion with EcoR1 and XbaI. A cDNA library was amplified from ~50,000 independent clones.

#### Isolation of Rat VEGF B cDNA Clone

- 10 [0108] The above cDNA library was screened by plaque hybridization using pCV2 as a probe. Hybridization conditions were as follows:

- 15 5XSSC ( 1XSSC is 0.15M sodium chloride, 0.015M sodium citrate,  
50% Formamide  
5X Denhardt's Solution ( 1% Ficoll, 1% polyvinylpyrrolidone, 1% bovine serum albumin)  
0.15 mg/ml salmon sperm DNA  
hybridize overnight at 42° C.

- 20 [0109] Filters were washed 3 times in 2XSSC, 0.1% SDS at room temperature for 5', then 1 time in 1XSSC, 0.1% SDS at 50C for 30'. Positive clones were identified by autoradiography.

- [0110] The DNA from phage #202 was digested with restriction endonuclease SpeI and the 1.1kb band ligated into XbaI digested pGEM3Zf(+). The ligation mix was used to transform *E.coli* XL-1 blue. Plasmid DNA was isolated from white transformants and sequenced by the dideoxy chain termination method. The cDNA sequence and predicted  
25 amino acid sequence of the signal peptide are shown in Figures 7 and 8.

- [0111] The entire nucleotide and amino acid sequence of the 138 amino acid form is shown in Fig. 7. The secreted protein starts at Ala<sup>24</sup> and continues to Arg<sup>138</sup>. The entire nucleotide and amino acid sequence of the 158 amino acid form is shown in Figures 8. The secreted protein starts at Ala<sup>24</sup> and continues to Leu<sup>158</sup>.

#### 30 EXAMPLE 11

##### Cloning and sequencing VEGF C Subunit

##### PCR Amplification, Cloning and Sequencing of pFSEM'

- 35 [0112] Two degenerate oligonucleotides were synthesized based on the sequence of rat VEGF B monomer in order to amplify VEGF cDNAs from the human medulloblastoma line TE-671, ATCC HTB (McAllister *et al.*, Int. J. Cancer 20: 206-212 [1977]). These oligonucleotides were:

- 40 FS 5' TTTGTCGACA TTC AGT CC(N) TC(N) TG(TC) GT 3' SEQ ID NO:21  
EM' 5' TTTGTCGACA CTG AGA GAA (N)GT CAT (CT)TC 3'  
SEQ ID NO:22  
where N= AGCT

- 45 [0113] Poly A+ RNA was isolated from TE-671 cells using the Fast Track RNA isolation kit from Invitrogen and the protocol provided. First strand cDNA synthesis was performed as follows using the cDNA Cycle kit from Invitrogen;

- 1µl 1µg of TE-671 polyA+ RNA  
19µl water  
50 5µl 100mM MeMgOH  
6.25µl 0.7M B-mercaptoethanol  
2.5µl random primer  
2.5µl RNase Inhibitor  
10µl 5X RT buffer  
55 2.5µl 25mM dNTPs  
1.25µl reverse transcriptase 12.5units

- [0114] The reaction was incubated for 60' at 42°C, then 3' at 95°C, placed on ice, then an additional 1.25ul reverse

transcriptase was added and the reaction incubated an additional 60' at 42°C.

[0115] The above procedure was performed in duplicate and the cDNAs pooled to a final volume of 100 $\mu$ l.

#### PCR Reactions:

5

#### Primary reaction (100 $\mu$ l)

[0116]

- |    |              |  |
|----|--------------|--|
| 10 | 10 $\mu$ l   | 10X buffer from Perkin Elmer Cetus GeneAmp kit |
|    | 16 $\mu$ l   | 1.25mM each of dATP, dCTP, dGTP, TTP           |
|    | 10 $\mu$ l   | first strand TE-671 cDNA                       |
|    | 2 $\mu$ l    | 50 pmoles FS primer                            |
|    | 2 $\mu$ l    | 50 pmoles EM' primer                           |
| 15 | 0.5 $\mu$ l  | 2.5 units Amplitaq DNA polymerase              |
|    | 59.5 $\mu$ l | water  |

[0117] Reaction conditions: 40 cycles of 90°C, 1'; 2' ramp to 45°C; 2' at 45°C; 2' at 72°C.

#### 20 Gel Purification

[0118] 20  $\mu$ l of the primary PCR reaction was purified on a 4% NuSieve agarose gel. The 180 base pair band was excised from the gel, heated to 65°C for 5' and used directly as template for the secondary PCR reaction.

#### 25 Secondary PCR reaction 200 $\mu$ l

[0119]

- |    |             |  |
|----|-------------|--|
|    | 20 $\mu$ l  | 10X buffer from Perkin Elmer Cetus GeneAmp kit |
| 30 | 32 $\mu$ l  | 1.25mM each of dATP, dCTP, dGTP, TTP           |
|    | 5 $\mu$ l   | melted gel slice                               |
|    | 4 $\mu$ l   | 100 pmoles FS primer                           |
|    | 4 $\mu$ l   | 100 pmoles EM' primer                          |
|    | 1 $\mu$ l   | 5 units Amplitaq DNA polymerase                |
| 35 | 134 $\mu$ l | water  |

[0120] Reaction conditions: 35 cycles of 94°C, 1'; 50°C, 2'; 72°C, 2'

[0121] The PCR product was purified on a Qiagen tip 5 column, then digested with restriction endonuclease Sall. The Sall fragment was then ligated into Sall cut pGEM3Z(+), and the ligation mix used to transform *E. coli* XL-1 blue. Plasmid DNA was isolated from white transformants and sequenced by the dideoxy chain termination method.

#### PCR Amplification, Cloning and Sequencing of p3'.19

[0122] Based on the sequence obtained from the pFSEM' clone, a specific PCR primer was synthesized; oligo LH 5' TTTGTGCGACA CTG CAC TGT GTG CCG GTG 3' SEQ ID NO:23. This primer was used in combination with oligo A17, 5' GACTCGAGTCGACATCG 3' SEQ ID NO:24, to amplify the cDNA encoding the COOH terminus of the VEGF C subunit using the 3' RACE technique described by Frohman et al., PNAS 85: 8998-9002 (1988).

[0123] Poly A+ RNA was isolated from TE-671 cells using the Fast Track RNA isolation kit from Invitrogen and the protocol provided. First strand cDNA synthesis was performed as follows using the cDNA Cycle kit from Invitrogen and the TA17 adapter primer:

50 TA17 5' GACTCGAGTCGACATCGATTTTTTTTTTTTTTTTTT 3' SEQ ID NO:5

- |    |              |                                |
|----|--------------|--------------------------------|
|    | 0.8 $\mu$ l  | 1 $\mu$ g of TE-671 polyA+ RNA |
|    | 20.7 $\mu$ l | water                          |
| 55 | 5 $\mu$ l    | 100 mM MeMgOH                  |
|    | 6.25 $\mu$ l | 0.7 M B-mercaptoethanol        |
|    | 1.0 $\mu$ l  | 0.88 $\mu$ g primer TA17       |
|    | 2.5 $\mu$ l  | RNase Inhibitor                |

10 µl 5X RT buffer  
 2.5 µl 25mM dNTPs  
 1.25 µl reverse transcriptase 12.5 units

- 5 [0124] The reaction was incubated for 60' at 42°C, then 3' at 95°C, placed on ice, then an additional 1.25ul reverse transcriptase was added and the reaction incubated an additional 60' at 42°C.

### 3' RACE PCR

#### 10 [0125]

20 µl 10 X buffer from Perkin Elmer Cetus GeneAmp kit  
 32 µl 1.25mM each of dATP, dCTP, dGTP, TTP  
 20 µl first strand TE-671 cDNA primed with TA17  
 15 2 µl 50 pmoles LH primer  
 2 µl 50 pmoles A17 primer  
 1.0 µl 5 units Amplitaq DNA polymerase  
 123 µl water

- 20 [0126] Reaction conditions: 40 cycles of 94C, 1'; 2' at 58°C; 3' at 72°C.

[0127] The PCR product was purified on a Qiagen tip 5 column, then digested with restriction endonuclease Sall. The Sall fragment was then ligated into Sall cut pGEM3Zf(+), and the ligation mix used to transform *E. coli* XL-1 blue. Plasmid DNA was isolated from white transformants and sequenced by the dideoxy chain termination method.

#### 25 PCR Amplification, Cloning and Sequencing of p5'.16

[0128] Based on the sequence obtained from the pFSEM' clone, two specific PCR primers were synthesized; oligo VE' 5' TTTGTCGACA AC ATT GGC CGT CTC CAC C 3' SEQ ID NO:24, and oligo TG' 5' TTTGTCGACA ATC GCC GCA GCA GCC GGT 3' SEQ ID NO:25. These primers were used in combination with

- 30 oligo A17, 5' GACTCGAGTCGACATCG 3' SEQ ID NO:8, and oligo TA17 5'GACTCGAGTCGACATCGATTTTTTTTTTTTTTTT 3' SEQ ID NO:5 to amplify the cDNA encoding the amino terminus of the VEGF C subunit using the 5' RACE technique described by Frohman et al., PNAS 85: 8998-9002 (1988).

- [0129] Poly A+ RNA was isolated from TE-671 cells using the Fast Track RNA isolation kit from Invitrogen and the protocol provided. First strand cDNA synthesis was performed as follows using the cDNA Cycle kit from Invitrogen and

35 the VE' primer:

1.0 µl 1 µg of TE-671 polyA+ RNA  
 20.25 µl water  
 5 µl 100 mM MeMgOH  
 40 6.25 µl 0.7 M B-mercaptoethanol  
 1.0 µl 1.0 µg primer VE'  
 2.5 µl RNase Inhibitor  
 10 µl 5X RT buffer  
 2.5 µl 25 mM dNTPs  
 45 0.5 µl AMV reverse transcriptase (Promega) 10units

- [0130] The reaction was incubated for 60' at 42°C, then 3' at 95°C, placed on ice, then an additional 1.25ul reverse transcriptase was added and the reaction incubated an additional 60' at 42°C. Excess oligo VE' was removed by a Centricon 100 spin column and the 5' end of the cDNA was tailed by the addition of dATP and terminal transferase. The

50 tailed cDNA was diluted to a final volume of 200 ul in 10mM Tris-HCl, 1mM EDTA, pH 7.5.

### 5' RACE PCR 5 X 100ul

#### [0131]

55 10 µl 10X buffer from Perkin Elmer Cetus GeneAmp kit  
 16 µl 1.25mM each of dATP, dCTP, dGTP, TTP  
 10 µl first strand TE-671 cDNA primed with VE'

2 µl 50 pmoles TG' primer  
 2 µl 50 pmoles A17 primer  
 2 µl 20 pmoles TA17 primer  
 0.5 µl 2.5 units Amplitaq DNA polymerase  
 5 57.5 µl water

[0132] Reaction conditions: 40 cycles of 94°C, 1'; 2' ramp to 58°C; 2' at 58°C; 2' at 72°C.

[0133] The PCR product was purified on a Qiagen tip 5 column, then digested with restriction endonuclease Sall. The Sall fragment was then ligated into Sall cut pGEM3Zf(+), and the ligation mix used to transform *E. coli* XL-1 blue. Plasmid DNA was isolated from white transformants and sequenced by the dideoxy chain termination method. The combined sequences form plasmids pFSEM', p3'19 and p5'16 are shown in Figure 9.

#### PCR Amplification, Cloning and Sequencing of phVC16 and phVC2

15 [0134] Based on the sequences of the p5'16 and p3'19 clones, two specific PCR primers were synthesized; oligo 5' GCVB 5' TTTGTCGAC TGG CTC TGG ACG TCT GAG 3' SEQ ID NO:26 and oligo 3'VC 5' TTTGTCGAC ACT GAA GAG TGT GAC GG 3' SEQ ID NO:27. These primers were used together to amplify the cDNA encoding the complete VEGF C subunit.

[0135] Poly A+ RNA was isolated from TE-671 cells using the Fast Track RNA isolation kit from Invitrogen and the protocol provided. First strand cDNA synthesis was performed as follows using the cDNA Cycle kit from Invitrogen;

0.8µl 1µg of TE-671 polyA+ RNA  
 19.2µl water  
 5µl 100 mM MeMgOH  
 25 6.25µl 0.7 M B-mercaptoethanol  
 2.5µl oligo dT primer  
 2.5µl RNase Inhibitor  
 10µl 5X RT buffer  
 2.5µl 25 mM dNTPs  
 30 1.25µl reverse transcriptase 12.5units

[0136] The reaction was incubated for 60' at 42C, then 3' at 95C, placed on ice, then an additional 1.25ul reverse transcriptase was added and the reaction incubated an additional 60' at 42C.

35 PCR Reaction 200 ul

#### [0137]

20µl 10X buffer from Perkin Elmer Cetus GeneAmp kit  
 40 32µl 1.25mM each of dATP, dCTP, dGTP, TTP  
 20µl first strand TE-671 cDNA primed with oligo dT  
 4µl 50 pmoles 5' GCVB primer  
 4µl 50 pmoles 3'VC primer  
 1µl 5 units Amplitaq DNA polymerase  
 45 119µl water

[0138] Reaction conditions: 40 cycles of 94°C, 1'; 2' at 50°C; 2' at 72°C.

[0139] The PCR product was purified on a Qiagen tip 5 column, then digested with restriction endonuclease Sall. The Sall fragment was then ligated into Sall cut pGEM3Zf(+), and the ligation mix used to transform *E. coli* XL-1 blue. Plasmid DNA was isolated from white transformants and sequenced by the dideoxy chain termination method. In the sequences of clones phVC16 and phVC2 base 463 (Fig. 9) was changed from a T to a C eliminating the translational stop codon following amino acid 154; this results in the addition of 16 amino acids following amino acid Lys 154. The nucleotide sequence and the deduced amino acid sequence of this addition is:

55

CAG AGA CCC ACA GAC TGC CAC CTG TGC GGC GAT GCT GTT  
 Gln Arg Pro Thr Asp Cys His Leu Cys Gly Asp Ala Val  
 155 160 165

CCC CGG AGG TAA  
 Pro Arg Arg

170

SEQ ID NO:29

[0140] In addition clone phVC16 contains a 3 base pair deletion (Figure 9, nucleotide residues 73-75) resulting in the deletion of Gln 25.

# SEQUENCE LISTING

- (2) INFORMATION FOR SEQ ID NO:1:  
 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 14 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: Not Applicable  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1  
 Ala Pro Thr Thr Glu Gly Glu Gln Lys Ala His Glu Val  
 5 10  
 Val

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: Not Applicable
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:  
 Ala Leu Ser Ala Gly Asn Xaa Ser Thr Glu Met Glu Val  
 5 10  
 Val Pro Phe Asn Glu Val  
 15

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:  
 TTTGTCGACT TYATGGAYGT NTAYCA 26

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 32 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:  
 CAGAGAATTC GTCGACARTC NGTRTTYTTR CA 32

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 35 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:  
 GACTCGAGTC GACATCGATT TTTTTTTTTT TTTT 35

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 28 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single

(D) TOPOLOGY: linear  
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:  
 TTTGTCGACA ACACAGGACG GCTTGAAG 28

(2) INFORMATION FOR SEQ ID NO:7:  
 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 25 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear  
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:  
 TTTGTCGACG AAAATCACTG TGAGC 25

(2) INFORMATION FOR SEQ ID NO:8:  
 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 17 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear  
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:  
 GACTCGAGTC GACATCG 17

(2) INFORMATION FOR SEQ ID NO:9:  
 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 28 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear  
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:  
 TTTGTCGACA ACACAGGACG GCTTGAAG 28

(2) INFORMATION FOR SEQ ID NO:10:  
 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 28 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear  
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:  
 TTTGTCGACA TACTCCTGGA AGATGTCC 28

(2) INFORMATION FOR SEQ ID NO:11:  
 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 18 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear  
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:  
 CTTTCATCATT GCAGCAGC 18



- 5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55
- (2) INFORMATION FOR SEQ ID NO:12:  
    (i) SEQUENCE CHARACTERISTICS:  
        (A) LENGTH: 26 base pairs  
        (B) TYPE: nucleic acid  
        (C) STRANDEDNESS: single  
        (D) TOPOLOGY: linear  
    (xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:  
        TTTGTCGACA ACCATGAACT TTCTGC 26
- (2) INFORMATION FOR SEQ ID NO:13:  
    (i) SEQUENCE CHARACTERISTICS:  
        (A) LENGTH: 26 base pairs  
        (B) TYPE: nucleic acid  
        (C) STRANDEDNESS: single  
        (D) TOPOLOGY: linear  
    (xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:  
        TTTGTCGACG GTGAGAGGTC TAGTTC 26
- (2) INFORMATION FOR SEQ ID NO:14:  
    (i) SEQUENCE CHARACTERISTICS:  
        (A) LENGTH: 26 base pairs  
        (B) TYPE: nucleic acid  
        (C) STRANDEDNESS: single  
        (D) TOPOLOGY: linear  
    (xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:  
        TTTGTCGACA TAYATHGCNG AYGARC 26
- (2) INFORMATION FOR SEQ ID NO:15:  
    (i) SEQUENCE CHARACTERISTICS:  
        (A) LENGTH: 26 base pairs  
        (B) TYPE: nucleic acid  
        (C) STRANDEDNESS: single  
        (D) TOPOLOGY: linear  
    (xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:  
        TTTGTCGACT CRTCRTRCA RCANCC 26
- (2) INFORMATION FOR SEQ ID NO:16:  
    (i) SEQUENCE CHARACTERISTICS:  
        (A) LENGTH: 27 base pairs  
        (B) TYPE: nucleic acid  
        (C) STRANDEDNESS: single  
        (D) TOPOLOGY: linear  
    (xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:  
        TTTGTCGACA CACCCTAATG AAGTGTC 27
- (2) INFORMATION FOR SEQ ID NO:17:  
    (i) SEQUENCE CHARACTERISTICS:  
        (A) LENGTH: 26 base pairs  
        (B) TYPE: nucleic acid  
        (C) STRANDEDNESS: single

(D) TOPOLOGY: linear  
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:  
TTTGTCGACA ACAGCGACTC AGAAGG 26

(2) INFORMATION FOR SEQ ID NO:18:  
(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 27 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear  
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:  
TTTGTCGACA CTGAATATAT GAGACAC 27

(2) INFORMATION FOR SEQ ID NO:19:  
(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 25 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear  
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:  
TTTGTCGACN NGCAGGTCCT AGCTG 25

(2) INFORMATION FOR SEQ ID NO:20:  
(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 26 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear  
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:  
TTTGTCGACN NCTAATAAAT AGAGGG 26

(2) INFORMATION FOR SEQ ID NO:21:  
(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 27 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear  
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:  
TTTGTCGACA TTCAGTCCNT CNTGYGT 27

(2) INFORMATION FOR SEQ ID NO:22:  
(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 28 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear  
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:  
TTTGTCGACA CTGAGAGAAN GTCATYTC 28

(2) INFORMATION FOR SEQ ID NO:23:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 28 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:  
 TTTGTGCGACA CTGCACTGTG TGCCGGTG 28

(2) INFORMATION FOR SEQ ID NO:24:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 28 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:  
 TTTGTGCGACA ACATTGGCCG TCTCCACC 28

(2) INFORMATION FOR SEQ ID NO:25:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 28 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:  
 TTTGTGCGACA ATCGCCGCAG CAGCCGGT 28

(2) INFORMATION FOR SEQ ID NO:26:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 27 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:  
 TTTGTGCGACT GGCTCTGGAC GTCTGAG

(2) INFORMATION FOR SEQ ID NO:27:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 26 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:  
 TTTGTGCGACA CTGAAGAGTG TGACGG 26

(2) INFORMATION FOR SEQ ID NO:28:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 51 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

5 CAG AGA CCC ACA GAC TGC CAC CTG TGC GGC GAT  
GCT GTT 39

CCC CGG AGG TAA  
51

(2) INFORMATION FOR SEQ ID NO:29:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 16

(B) TYPE: amino acids

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

20 Gln Arg Pro Thr Asp Cys His Leu Cys Gly Asp  
Ala Val

5

10

Pro Arg Arg  
15

(2) INFORMATION FOR SEQ ID NO:30:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 577 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

AACC

4

35 ATG AAC TTT CTG CTC TCT TGG GTG CAC TGG ACC CTG GCT TTA CTG 49  
Met Asn Phe Leu Leu Ser Trp Val His Trp Thr Leu Ala Leu Leu  
5 10 15

40 CTG TAC CTC CAC CAT GCC AAG TGG TCC CAG GCT GCA CCC ACG ACA 94  
Leu Tyr Leu His His Ala Lys Trp Ser Gln Ala Ala Pro Thr Thr  
20 25 30

45 GAA GGG GAG CAG AAA GCC CAT GAA GTG GTG AAG TTC ATG GAC GTC 139  
Glu Gly Glu Gln Lys Ala His Glu Val Val Lys Phe Met Asp Val  
35 40 40

TAC CAG CGC AGC TAT TGC CGT CCG ATT GAG ACC CTG GTG GAC ATC 184  
Tyr Gln Arg Ser Tyr Cys Arg Pro Ile Glu Thr Leu Val Asp Ile  
50 55 60

55 TTC CAG GAG TAC CCC GAT GAG ATA GAG TAT ATC TTC AAG CCG TCC 229  
Phe Gln Glu Tyr Pro Asp Glu Ile Glu Tyr Ile Phe Lys Pro Ser  
65 70 75

5 TGT GTG CCC CTA ATG CGG TGT GCG GGC TGC TGC AAT GAT GAA GCC 274  
 Cys Val Pro Leu Met Arg Cys Ala Gly Cys Cys Asn Asp Glu Ala  
 80 85 90

CTG GAG TGC GTG CCC ACG TCG GAG AGC AAC GTC ACT ATG CAG ATC 319  
 Leu Glu Cys Val Pro Thr Ser Glu Ser Asn Val Thr Met Gln Ile  
 95 100 105

10 ATG CGG ATC AAA CCT CAC CAA AGC CAG CAC ATA GGA GAG ATG AGC 364  
 Met Arg Ile Lys Pro His Gln Ser Gln His Ile Gly Glu Met Ser  
 110 115 120

15 TTC CTG CAG CAT AGC AGA TGT GAA TGC AGA CCA AAG AAA GAT AGA 409  
 Phe Leu Gln His Ser Arg Cys Glu Cys Arg Pro Lys Lys Asp Arg  
 125 130 135

20 ACA AAG CCA GAA AAT CAC TGT GAG CCT TGT TCA GAG CGG AGA AAG 454  
 Thr Lys Pro Glu Asn His Cys Glu Pro Cys Ser Glu Arg Arg Lys  
 140 145 150

CAT TTG TTT GTC CAA GAT CCG CAG ACG TGT AAA TGT TCC TGC AAA 499  
 His Leu Phe Val Gln Asp Pro Gln Thr Cys Lys Cys Ser Cys Lys  
 155 160 165

25 AAC ACA GAC TCG CGT TGC AAG GCG AGG CAG CTT GAG TTA AAC GAA 544  
 Asn Thr Asp Ser Arg Cys Lys Ala Arg Gln Leu Glu Leu Asn Glu  
 170 175 180

30 CGT ACT TGC AGA TGT GAC AAG CCA AGG CGG TGA 577  
 Arg Thr Cys Arg Cys Asp Lys Pro Arg Arg  
 185 190

## (2) INFORMATION FOR SEQ ID NO:31:

## (i) SEQUENCE CHARACTERISTICS:

- 35 (A) LENGTH: 190 amino acids  
 (B) TYPE: amino acids  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

40 Met Asn Phe Leu Leu Ser Trp Val His Trp Thr Leu Ala Leu Leu  
 5 10 15

Leu Tyr Leu His His Ala Lys Trp Ser Gln Ala Ala Pro Thr Thr  
 20 25 30

45 Glu Gly Glu Gln Lys Ala His Glu Val Val Lys Phe Met Asp Val  
 35 40 45

50 Tyr Gln Arg Ser Tyr Cys Arg Pro Ile Glu Thr Leu Val Asp Ile  
 50 55 60

55 Phe Gln Glu Tyr Pro Asp Glu Ile Glu Tyr Ile Phe Lys Pro Ser

	65	70	75
5	Cys Val Pro Leu Met Arg Cys Ala Gly Cys Cys Asn Asp Glu Ala 80	85	90
	Leu Glu Cys Val Pro Thr Ser Glu Ser Asn Val Thr Met Gln Ile 95	100	105
10	Met Arg Ile Lys Pro His Gln Ser Gln His Ile Gly Glu Met Ser 110	115	120
	Phe Leu Gln His Ser Arg Cys Glu Cys Arg Pro Lys Lys Asp Arg 125	130	135
15	Thr Lys Pro Glu Asn His Cys Glu Pro Cys Ser Glu Arg Arg Lys 140	145	150
	His Leu Phe Val Gln Asp Pro Gln Thr Cys Lys Cys Ser Cys Lys 155	160	165
20	Asn Thr Asp Ser Arg Cys Lys Ala Arg Gln Leu Glu Leu Asn Glu 170	175	180
	Arg Thr Cys Arg Cys Asp Lys Pro Arg Arg 185	190	

## (2) INFORMATION FOR SEQ ID NO:32:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 445 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

35	ACCA	4
	ATG AAC TTT CTG CTC TCT TGG GTG CAC TGG ACC CTG GCT TTA CTG Met Asn Phe Leu Leu Ser Trp Val His Trp Thr Leu Ala Leu Leu 5 10 15	49
40	CTG TAC CTC CAC CAT GCC AAG TGG TCC CAG GCT GCA CCC ACG ACA Leu Tyr Leu His His Ala Lys Trp Ser Gln Ala Ala Pro Thr Thr 20 25 30	94
45	GAA GGG GAG CAG AAA GCC CAT GAA GTG GTG AAG TTC ATG GAC GTC Glu Gly Glu Gln Lys Ala His Glu Val Val Lys Phe Met Asp Val 35 40 45	139
50	TAC CAG CGC AGC TAT TGC CGT CCG ATT GAG ACC CTG GTG GAC ATC Tyr Gln Arg Ser Tyr Cys Arg Pro Ile Glu Thr Leu Val Asp Ile 50 55 60	184
55	TTC CAG GAG TAC CCC GAT GAG ATA GAG TAT ATC TTC AAG CCG TCC Phe Gln Glu Tyr Pro Asp Glu Ile Glu Tyr Ile Phe Lys Pro Ser	229

	65	70	75	
5	TGT GTG CCC CTA ATG CGG TGT GCG GGC TGC TGC AAT GAT GAA GCC	274		
	Cys Val Pro Leu Met Arg Cys Ala Gly Cys Cys Asn Asp Glu Ala			
	80	85	90	
10	CTG GAG TGC GTG CCC ACG TCG GAG AGC AAC GTC ACT ATG CAG ATC	319		
	Leu Glu Cys Val Pro Thr Ser Glu Ser Asn Val Thr Met Gln Ile			
	95	100	105	
15	ATG CGG ATC AAA CCT CAC CAA AGC CAG CAC ATA GGA GAG ATG AGC	364		
	Met Arg Ile Lys Pro His Gln Ser Gln His Ile Gly Glu Met Ser			
	110	115	120	
20	TTC CTG CAG CAT AGC AGA TGT GAA TGC AGA CCA AAG AAA GAT AGA	409		
	Phe Leu Gln His Ser Arg Cys Glu Cys Arg Pro Lys Lys Asp Arg			
	125	130	135	
25	ACA AAG CCA GAA AAA TGT GAC AAG CCA AGG CGG TGA	445		
	Thr Lys Pro Glu Lys Cys Asp Lys Pro Arg Arg			
	140	145		

## (2) INFORMATION FOR SEQ ID NO:33:

## (i) SEQUENCE CHARACTERISTICS:

- 25 (A) LENGTH: 146 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

30	Met Asn Phe Leu Leu Ser Trp Val His Trp Thr Leu Ala Leu Leu	
	5	10
35	Leu Tyr Leu His His Ala Lys Trp Ser Gln Ala Ala Pro Thr Thr	
	20	25
40	Glu Gly Glu Gln Lys Ala His Glu Val Val Lys Phe Met Asp Val	
	35	40
45	Tyr Gln Arg Ser Tyr Cys Arg Pro Ile Glu Thr Leu Val Asp Ile	
	50	55
50	Phe Gln Glu Tyr Pro Asp Glu Ile Glu Tyr Ile Phe Lys Pro Ser	
	65	70
55	Cys Val Pro Leu Met Arg Cys Ala Gly Cys Cys Asn Asp Glu Ala	
	80	85
60	Leu Glu Cys Val Pro Thr Ser Glu Ser Asn Val Thr Met Gln Ile	
	95	100
65	Met Arg Ile Lys Pro His Gln Ser Gln His Ile Gly Glu Met Ser	
	110	115
70		120

Phe Leu Gln His Ser Arg Cys Glu Cys Arg Pro Lys Lys Asp Arg  
125 130 135

Thr Lys Pro Glu Lys Cys Asp Lys Pro Arg Arg  
140 145

## (2) INFORMATION FOR SEQ ID NO:34:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 649 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ  
ID NO:34:

AACC

4

ATG AAC TTT CTG CTC TCT TGG GTG CAC TGG ACC CTG GCT TTA CTG 49  
Met Asn Phe Leu Leu Ser Trp Val His Trp Thr Leu Ala Leu Leu  
5 10 15

CTG TAC CTC CAC CAT GCC AAG TGG TCC CAG GCT GCA CCC ACG ACA 94  
Leu Tyr Leu His His Ala Lys Trp Ser Gln Ala Ala Pro Thr Thr  
20 25 30

GAA GGG GAG CAG AAA GCC CAT GAA GTG GTG AAG TTC ATG GAC GTC 139  
Glu Gly Glu Gln Lys Ala His Glu Val Val Lys Phe Met Asp Val  
35 40 45

TAC CAG CGC AGC TAT TGC CGT CCG ATT GAG ACC CTG GTG GAC ATC 184  
Tyr Gln Arg Ser Tyr Cys Arg Pro Ile Glu Thr Leu Val Asp Ile  
50 55 60

TTC CAG GAG TAC CCC GAT GAG ATA GAG TAT ATC TTC AAG CCG TCC 229  
Phe Gln Glu Tyr Pro Asp Glu Ile Glu Tyr Ile Phe Lys Pro Ser  
65 70 75

TGT GTG CCC CTA ATG CGG TGT GCG GGC TGC TGC AAT GAT GAA GCC 274  
Cys Val Pro Leu Met Arg Cys Ala Gly Cys Cys Asn Asp Glu Ala  
80 85 90

CTG GAG TGC GTG CCC ACG TCG GAG AGC AAC GTC ACT ATG CAG ATC 319  
Leu Glu Cys Val Pro Thr Ser Glu Ser Asn Val Thr Met Gln Ile  
95 100 105

ATG CGG ATC AAA CCT CAC CAA AGC CAG CAC ATA GGA GAG ATG AGC 364  
Met Arg Ile Lys Pro His Gln Ser Gln His Ile Gly Glu Met Ser  
110 115 120

TTC CTG CAG CAT AGC AGA TGT GAA TGC AGA CCA AAG AAA GAT AGA 409  
Phe Leu Gln His Ser Arg Cys Glu Cys Arg Pro Lys Lys Asp Arg  
125 130 135

ACA AAG CCA GAA AAA AAA TCA GTT CGA GGA AAG GGA AAG GGT CAA 454



Thr Lys Pro Glu Lys Lys Ser Val Arg Gly Lys Gly Lys Gly Gln  
 140 145 150  
 5 AAA CGA AAG CGC AAG AAA TCC CGG TTT AAA TCC TGG AGC GTT CAC 499  
 Lys Arg Lys Arg Lys Lys Ser Arg Phe Lys Ser Trp Ser Val His  
 155 160 165  
 TGT GAG CCT TGT TCA GAG CGG AGA AAG CAT TTG TTT GTC CAA GAT 544  
 10 Cys Glu Pro Cys Ser Glu Arg Arg Lys His Leu Phe Val Gln Asp  
 170 175 180  
 CCG CAG ACG TGT AAA TGT TCC TGC AAA AAC ACA GAC TCG CGT TGC 589  
 15 Pro Gln Thr Cys Lys Cys Ser Cys Lys Asn Thr Asp Ser Arg Cys  
 185 190 195  
 AAG GCG AGG CAG CTT GAG TTA AAC GAA CGT ACT TGC AGA TGT GAC 634  
 Lys Ala Arg Gln Leu Glu Leu Asn Glu Arg Thr Cys Arg Cys Asp  
 200 205 210  
 20 AAG CCA AGG CGG TGA 649  
 Lys Pro Arg Arg

## (2) INFORMATION FOR SEQ ID NO:35:

## (i) SEQUENCE CHARACTERISTICS:

- 25 (A) LENGTH: 214 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

30 Met Asn Phe Leu Leu Ser Trp Val His Trp Thr Leu Ala Leu Leu  
 5 10 15  
 Leu Tyr Leu His His Ala Lys Trp Ser Gln Ala Ala Pro Thr Thr  
 35 20 25 30  
 Glu Gly Glu Gln Lys Ala His Glu Val Val Lys Phe Met Asp Val  
 35 40 45  
 Tyr Gln Arg Ser Tyr Cys Arg Pro Ile Glu Thr Leu Val Asp Ile  
 40 50 55 60  
 Phe Gln Glu Tyr Pro Asp Glu Ile Glu Tyr Ile Phe Lys Pro Ser  
 65 70 75  
 45 Cys Val Pro Leu Met Arg Cys Ala Gly Cys Cys Asn Asp Glu Ala  
 80 85 90  
 Leu Glu Cys Val Pro Thr Ser Glu Ser Asn Val Thr Met Gln Ile  
 95 100 105  
 50 Met Arg Ile Lys Pro His Gln Ser Gln His Ile Gly Glu Met Ser  
 110 115 120

55

Phe Leu Gln His Ser Arg Cys Glu Cys Arg Pro Lys Lys Asp Arg  
125 130 135

Thr Lys Pro Glu Lys Lys Ser Val Arg Gly Lys Gly Lys Gly Gln  
140 145 150

Lys Arg Lys Arg Lys Lys Ser Arg Phe Lys Ser Trp Ser Val His  
155 160 165

Cys Glu Pro Cys Ser Glu Arg Arg Lys His Leu Phe Val Gln Asp  
170 175 180

Pro Gln Thr Cys Lys Cys Ser Cys Lys Asn Thr Asp Ser Arg Cys  
185 190 195

Lys Ala Arg Gln Leu Glu Leu Asn Glu Arg Thr Cys Arg Cys Asp  
200 205 210

Lys Pro Arg Arg

(2) INFORMATION FOR SEQ ID NO:36:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 417 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

ATG CTG GCC ATG AAG CTG TTC ACT TGC TTC TTG CAG GTC CTA GCT 45  
Met Leu Ala Met Lys Leu Phe Thr Cys Phe Leu Gln Val Leu Ala  
5 10 15

GGG TTG GCT GTG CAC TCC CAG GGG GCC CTG TCT GCT GGG AAC AAC 90  
Gly Leu Ala Val His Ser Gln Gly Ala Leu Ser Ala Gly Asn Asn  
20 25 30

TCA ACA GAA ATG GAA GTG GTG CCT TTC AAT GAA GTG TGG GGC CGC 135  
Ser Thr Glu Met Glu Val Val Pro Phe Asn Glu Val Trp Gly Arg  
35 40 45

AGC TAC TGC CGG CCA ATG GAG AAG CTG GTG TAC ATT GCA GAT GAA 180  
Ser Tyr Cys Arg Pro Met Glu Lys Leu Val Tyr Ile Ala Asp Glu  
50 55 60

CAC CCT AAT GAA GTG TCT CAT ATA TTC AGT CCG TCA TGT GTC CTT 225  
His Pro Asn Glu Val Ser His Ile Phe Ser Pro Ser Cys Val Leu  
65 70 75

CTG AGT CGC TGT AGT GGC TGC TGT GGT GAC GAG GGT CTG CAC TGT 270  
Leu Ser Arg Cys Ser Gly Cys Cys Gly Asp Glu Gly Leu His Cys  
80 85 90

GTG GCG CTA AAG ACA GCC AAC ATC ACT ATG CAG ATC TTA AAG ATT 315  
 Val Ala Leu Lys Thr Ala Asn Ile Thr Met Gln Ile Leu Lys Ile  
                             95                            100                            105

CCC CCC AAT CGG GAT CCA CAT TCC TAC GTG GAG ATG ACA TTC TCT 360  
 Pro Pro Asn Arg Asp Pro His Ser Tyr Val Glu Met Thr Phe Ser  
                             110                            115                            120

CAG GAT GTA CTC TGC GAA TGC AGG CCT ATT CTG GAG ACG ACA AAG 405  
 Gln Asp Val Leu Cys Glu Cys Arg Pro Ile Leu Glu Thr Thr Lys  
                             125                            130                            135

GCA GAA AGG TAA 417  
 Ala Glu Arg

(2) INFORMATION FOR SEQ ID NO:37:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 138 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

Met Leu Ala Met Lys Leu Phe Thr Cys Phe Leu Gln Val Leu Ala  
                             5                            10                            15

Gly Leu Ala Val His Ser Gln Gly Ala Leu Ser Ala Gly Asn Asn  
                             20                            25                            30

Ser Thr Glu Met Glu Val Val Pro Phe Asn Glu Val Trp Gly Arg  
                             35                            40                            45

Ser Tyr Cys Arg Pro Met Glu Lys Leu Val Tyr Ile Ala Asp Glu  
                             50                            55                            60

His Pro Asn Glu Val Ser His Ile Phe Ser Pro Ser Cys Val Leu  
                             65                            70                            75

Leu Ser Arg Cys Ser Gly Cys Cys Gly Asp Glu Gly Leu His Cys  
                             80                            85                            90

Val Ala Leu Lys Thr Ala Asn Ile Thr Met Gln Ile Leu Lys Ile  
                             95                            100                            105

Pro Pro Asn Arg Asp Pro His Ser Tyr Val Glu Met Thr Phe Ser  
                             110                            115                            120

Gln Asp Val Leu Cys Glu Cys Arg Pro Ile Leu Glu Thr Thr Lys  
                             125                            130                            135

Ala Glu Arg

(2) INFORMATION FOR SEQ ID NO:38:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 477 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

ATG CTG GCC ATG AAG CTG TTC ACT TGC TTC TTG CAG GTC CTA GCT 45  
 Met Leu Ala Met Lys Leu Phe Thr Cys Phe Leu Gln Val Leu Ala  
                   5                  10                  15

GGG TTG GCT GTG CAC TCC CAG GGG GCC CTG TCT GCT GGG AAC AAC 90  
 Gly Leu Ala Val His Ser Gln Gly Ala Leu Ser Ala Gly Asn Asn  
                   20                  25                  30

TCA ACA GAA ATG GAA GTG GTG CCT TTC AAT GAA GTG TGG GGC CGC 135  
 Ser Thr Glu Met Glu Val Val Pro Phe Asn Glu Val Trp Gly Arg  
                   35                  40                  45

AGC TAC TGC CGG CCA ATG GAG AAG CTG GTG TAC ATT GCA GAT GAA 180  
 Ser Tyr Cys Arg Pro Met Glu Lys Leu Val Tyr Ile Ala Asp Glu  
                   50                  55                  60

CAC CCT AAT GAA GTG TCT CAT ATA TTC AGT CCG TCA TGT GTC CTT 225  
 His Pro Asn Glu Val Ser His Ile Phe Ser Pro Ser Cys Val Leu  
                   65                  70                  75

CTG AGT CGC TGT AGT GGC TGC TGT GGT GAC GAG GGT CTG CAC TGT 270  
 Leu Ser Arg Cys Ser Gly Cys Cys Gly Asp Glu Gly Leu His Cys  
                   80                  85                  90

GTG GCG CTA AAG ACA GCC AAC ATC ACT ATG CAG ATC TTA AAG ATT 315  
 Val Ala Leu Lys Thr Ala Asn Ile Thr Met Gln Ile Leu Lys Ile  
                   95                  100                  105

CCC CCC AAT CGG GAT CCA CAT TCC TAC GTG GAG ATG ACA TTC TCT 360  
 Pro Pro Asn Arg Asp Pro His Ser Tyr Val Glu Met Thr Phe Ser  
                   110                  115                  120

CAG GAT GTA CTC TGC GAA TGC AGG CCT ATT CTG GAG ACG ACA AAG 405  
 Gln Asp Val Leu Cys Glu Cys Arg Pro Ile Leu Glu Thr Thr Lys  
                   125                  130                  135

GCA GAA AGG AGG AAA ACC AAG GGG AAG AGG AAG CAA AGC AAA ACC 450  
 Ala Glu Arg Arg Lys Thr Lys Gly Lys Arg Lys Gln Ser Lys Thr  
                   140                  145                  150

CCA CAG ACT GAG GAA CCC CAC CTG TGA 477  
 Pro Gln Thr Glu Glu Pro His Leu  
                   155

(2) INFORMATION FOR SEQ ID NO:39:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 158 amino acids

(B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:

Met Leu Ala Met Lys Leu Phe Thr Cys Phe Leu Gln Val Leu Ala 15  
                                   5                                  10  
 10 Gly Leu Ala Val His Ser Gln Gly Ala Leu Ser Ala Gly Asn Asn  
                                   20                                  25                                  30  
 Ser Thr Glu Met Glu Val Val Pro Phe Asn Glu Val Trp Gly Arg 45  
 15                                  35                                  40  
 Ser Tyr Cys Arg Pro Met Glu Lys Leu Val Tyr Ile Ala Asp Glu 60  
                                   50                                  55  
 20 His Pro Asn Glu Val Ser His Ile Phe Ser Pro Ser Cys Val Leu 75  
                                   65                                  70  
 Leu Ser Arg Cys Ser Gly Cys Cys Gly Asp Glu Gly Leu His Cys 90  
                                   80                                  85  
 25 Val Ala Leu Lys Thr Ala Asn Ile Thr Met Gln Ile Leu Lys Ile 105  
                                   95                                  100  
 Pro Pro Asn Arg Asp Pro His Ser Tyr Val Glu Met Thr Phe Ser 120  
                                   110                                  115  
 30 Gln Asp Val Leu Cys Glu Cys Arg Pro Ile Leu Glu Thr Thr Lys 135  
                                   125                                  130  
 Ala Glu Arg Arg Lys Thr Lys Gly Lys Arg Lys Gln Ser Lys Thr 150  
 35                                  140                                  145  
 Pro Gln Thr Glu Glu Pro His Leu 155

(2) INFORMATION FOR SEQ ID NO:40:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 465 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:

ATG CCG GTC ATG AGG CTG TTC CCT TGC TTC CTG CAG CTC CTG GCC 45  
 Met Pro Val Met Arg Leu Phe Pro Cys Phe Leu Gln Leu Leu Ala  
                                   5                                  10                                  15

50

55

GGG CTG GCG CTG CCT GCT GTG CCC CCC CAG CAG TGG GCC TTG TCT 90  
 Gly Leu Ala Leu Pro Ala Val Pro Pro Gln Gln Trp Ala Leu Ser  
                             20                            25                            30

GCT GGG AAC GGC TCG TCA GAG GTG GAA GTG GTA CCC TTC CAG GAA 135  
 Ala Gly Asn Gly Ser Ser Glu Val Glu Val Val Pro Phe Gln Glu  
                             35                            40                            45

GTG TGG GGC CGC AGC TAC TGC CGG GCG CTG GAG AGG CTG GTG GAC 180  
 Val Trp Gly Arg Ser Tyr Cys Arg Ala Leu Glu Arg Leu Val Asp  
                             50                            55                            60

GTC GTG TCC GAG TAC CCC AGC GAG GTG GAG CAC ATG TTC AGC CCA 225  
 Val Val Ser Glu Tyr Pro Ser Glu Val Glu His Met Phe Ser Pro  
                             65                            70                            75

TCC TGT GTC TCC CTG CTG CGC TGC ACC GGC TGC TGC GGC GAT GAG 270  
 Ser Cys Val Ser Leu Leu Arg Cys Thr Gly Cys Cys Gly Asp Glu  
                             80                            85                            90

AAT CTG CAC TGT GTG CCG GTG GAG ACG GCC AAT GTC ACC ATG CAG 315  
 Asn Leu His Cys Val Pro Val Glu Thr Ala Asn Val Thr Met Gln  
                             95                            100                            105

CTC CTA AAG ATC CGT TCT GGG GAC CGG CCC TCC TAC GTG GAG CTG 360  
 Leu Leu Lys Ile Arg Ser Gly Asp Arg Pro Ser Tyr Val Glu Leu  
                             110                            115                            120

ACG TTC TCT CAG CAC GTT CGC TGC GAA TGC CGG CCT CTG CGG GAG 405  
 Thr Phe Ser Gln His Val Arg Cys Glu Cys Arg Pro Leu Arg Glu  
                             125                            130                            135

AAG ATG AAG CCG GAA AGG AGG AGA CCC AAG GGC AGG GGG AAG AGG 450  
 Lys Met Lys Pro Glu Arg Arg Arg Pro Lys Gly Arg Gly Lys Arg  
                             140                            145                            150

AGG AGA GAG AAG TAG 465  
 Arg Arg Glu Lys

(2) INFORMATION FOR SEQ ID NO:41:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 154 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

Met Pro Val Met Arg Leu Phe Pro Cys Phe Leu Gln Leu Leu Ala  
                             5                            10                            15

Gly Leu Ala Leu Pro Ala Val Pro Pro Gln Gln Trp Ala Leu Ser  
                             20                            25                            30

Ala Gly Asn Gly Ser Ser Glu Val Glu Val Val Pro Phe Gln Glu  
 35 40 45  
 5 Val Trp Gly Arg Ser Tyr Cys Arg Ala Leu Glu Arg Leu Val Asp  
 50 55 60  
 Val Val Ser Glu Tyr Pro Ser Glu Val Glu His Met Phe Ser Pro  
 65 70 75  
 10 Ser Cys Val Ser Leu Leu Arg Cys Thr Gly Cys Cys Gly Asp Glu  
 80 85 90  
 Asn Leu His Cys Val Pro Val Glu Thr Ala Asn Val Thr Met Gln  
 95 100 105  
 15 Leu Leu Lys Ile Arg Ser Gly Asp Arg Pro Ser Tyr Val Glu Leu  
 110 115 120  
 Thr Phe Ser Gln His Val Arg Cys Glu Cys Arg Pro Leu Arg Glu  
 125 130 135  
 20 Lys Met Lys Pro Glu Arg Arg Arg Pro Lys Gly Arg Gly Lys Arg  
 140 145 150  
 25 Arg Arg Glu Lys

## 30 Claims

1. A purified and isolated DNA sequence encoding the C subunit of vascular endothelial cell growth factor comprising:

35

40

45

50

55

5 ATG CCG GTC ATG AGG CTG TTC OCT TGC TTC CTG CAG CTC CTG GOC  
 GGG CTG GCG CTG OCT GCT GTG CCC CCC CAG CAG TGG GOC TTG TCT  
 GCT GGG AAC GGC TCG TCA GAG GTG GAA GTG GTA CCC TTC CAG GAA  
 10 GTG TGG GGC CGC AGC TAC TGC CCG GCG CTG GAG AGG CTG GTG GAC  
 GTC GTG TCC GAG TAC CCC AGC GAG GTG GAG CAC ATG TTC AGC CCA  
 15 TCC TGT GTC TCC CTG CTG CGC TGC ACC GGC TGC TGC GGC GAT GAG  
 AAT CTG CAC TGT GTG CCG GTG GAG ACG GOC AAT GTC ACC ATG CAG  
 CTC CTA AAG ATC CGT TCT GGG GAC CCG CCC TCC TAC GTG GAG CTG  
 20 ACG TTC TCT CAG CAC GTT CGC TGC GAA TGC CCG OCT CTG CCG GAG  
 AAG ATG AAG CCG GAA AGG AGG AGA CCC AAG GGC AGG GGG AAG AGG  
 25 AGG AGA GAG AAG TAG

- 30 2. A purified and isolated vascular endothelial cell growth factor C subunit DNA sequence comprising:

ATG CCG GTC ATG AGG CTG TTC OCT TGC TTC CTG CAG CTC CTG  
 35 GOC GGG CTG GCG CTG OCT GCT GTG CCC CCC CAG CAG TGG GCC  
 TTG TCT GCT GGG AAC GGC TCG TCA GAG GTG GAA GTG GTA CCC  
 TTC CAG GAA GTG TGG GGC CGC AGC TAC TGC CCG GCG CTG GAG  
 40 AGG CTG GTG GAC GTC GTG TCC GAG TAC CCC AGC GAG GTG GAG  
 CAC ATG TTC AGC CCA TCC TGT GTC TCC CTG CTG CGC TGC ACC  
 GGC TGC TGC GGC GAT GAG AAT CTG CAC TGT GTG CCG GTG GAG  
 ACG GOC AAT GTC ACC ATG CAG CTC CTA AAG ATC CGT TCT GGG  
 45 GAC CCG CCC TCC TAC GTG GAG CTG ACG TTC TCT CAG CAC GTT  
 CGC TGC GAA TGC CCG OCT CTG CCG GAG AAG ATG AAG CCG GAA  
 AGG AGG AGA CCC AAG GGC AGG GGG AAG AGG AGA GAG AAG  
 50 CAG AGA CCC ACA GAC TGC CAC CTG TGC GGC GAT GCT GTT CCC  
 CCG AGG TAA. SEQ ID NOS: 28 & 40

- 55 3. Vascular endothelial cell growth factor AC DNA comprising an A subunit DNA sequence as depicted in Figure 4, 5 or 6 herein and a C subunit DNA sequence as claimed in claim 1 or 2.



4. Vascular endothelial cell growth factor BC DNA comprising a B subunit DNA sequence as depicted in Figure 7 or 8 herein and a C subunit DNA sequence as claimed in claim 1 or 2.
5. Vascular endothelial cell growth factor AC DNA comprising an A subunit DNA sequence selected from the group consisting of: a DNA sequence encoding a 189 amino acid form, a DNA sequence encoding a 165 amino acid form and a DNA sequence encoding a 121 amino acid form, with said A subunit DNA operably attached to a C subunit DNA sequence as claimed in claim 1 or 2.
6. Vascular endothelial cell growth factor BC DNA comprising a B subunit DNA sequence selected from the group consisting of a DNA sequence encoding a 135 amino acid form as depicted in Figure 8 herein starting at Ala<sup>24</sup> and continuing to Leu<sup>158</sup>, and a DNA sequence encoding a 115 amino acid form as depicted in Figure 7 herein starting at Ala<sup>24</sup> and continuing to Arg<sup>138</sup>, with said B subunit DNA sequence operably attached to a C subunit DNA sequence as claimed in claim 1 or 2.
7. Homodimeric vascular endothelial growth factor DNA comprising C subunit DNA sequences as claimed in claim 1 or 2.
8. A vector containing the DNA sequence of any one of claims 2 to 7.
9. A host cell transformed by the vector of claim 8 containing the DNA sequence encoding vascular endothelial cell growth factor.
10. A process for the preparation of vascular endothelial cell growth factor comprising culturing the transformed host cell of claim 9 under conditions suitable for the expression of vascular endothelial cell growth factor and recovering vascular endothelial cell growth factor.
11. Vascular endothelial growth factor made by the process of claim 10.
12. A purified and isolated vascular endothelial cell growth factor C subunit amino acid sequence comprising:

Met Pro Val Met Arg Leu Phe Pro Cys Phe Leu Gln Leu Leu  
 Ala Gly Leu Ala Leu Pro Ala Val Pro Pro Gln Gln Trp Ala  
 Leu Ser Ala Gly Asn Gly Ser Ser Glu Val Glu Val Val Pro  
 Phe Gln Glu Val Trp Gly Arg Ser Tyr Cys Arg Ala Leu Glu  
 Arg Leu Val Asp Val Val Ser Glu Tyr Pro Ser Glu Val Glu  
 His Met Phe Ser Pro Ser Cys Val Ser Leu Leu Arg Cys Thr  
 Gly Cys Cys Gly Asp Glu Asn Leu His Cys Val Pro Val Glu  
 Thr Ala Asn Val Thr Met Gln Leu Leu Lys Ile Arg Ser Gly  
 Asp Arg Pro Ser Tyr Val Glu Leu Thr Phe Ser Gln His Val  
 Arg Cys Glu Cys Arg Pro Leu Arg Glu Lys Met Lys Pro Glu  
 Arg Arg Arg Pro Lys Gly Arg Gly Lys Arg Arg Arg Glu Lys.  
 SEQ ID NO:41

13. A purified and isolated vascular endothelial cell growth factor C subunit amino acid sequence comprising:

Met Pro Val Met Arg Leu Phe Pro Cys Phe Leu Gln Leu Leu  
 Ala Gly Leu Ala Leu Pro Ala Val Pro Pro Gln Gln Trp Ala  
 5 Leu Ser Ala Gly Asn Gly Ser Ser Glu Val Glu Val Val Pro  
 Phe Gln Glu Val Trp Gly Arg Ser Tyr Cys Arg Ala Leu Glu  
 Arg Leu Val Asp Val Val Ser Glu Tyr Pro Ser Glu Val Glu  
 10 His Met Phe Ser Pro Ser Cys Val Ser Leu Leu Arg Cys Thr  
 Gly Cys Cys Gly Asp Glu Asn Leu His Cys Val Pro Val Glu  
 Thr Ala Asn Val Thr Met Gln Leu Leu Lys Ile Arg Ser Gly  
 15 Asp Arg Pro Ser Tyr Val Glu Leu Thr Phe Ser Gln His Val  
 Arg Cys Glu Cys Arg Pro Leu Arg Glu Lys Met Lys Pro Glu  
 Arg Arg Arg Pro Lys Gly Arg Gly Lys Arg Arg Arg Glu Lys  
 20 Gln Arg Pro Thr Asp Cys His Leu Cys Gly Asp Ala Val Pro  
 Arg Arg. SEQ ID NOS: 29 & 40

- 25
14. Vascular endothelial cell growth factor AC comprising an A subunit amino acid sequence as depicted in Figure 4, 5 or 6 herein and a C subunit amino acid sequence as claimed in claim 12 or 13.
- 30
15. Vascular endothelial cell growth factor BC comprising a B subunit amino acid sequence as depicted in Figure 7 or 8 herein and a C subunit amino acid sequence as claimed in claim 12 or 13.
- 35
16. Vascular endothelial cell growth factor CC comprising a first C subunit amino acid sequence as claimed in claim 12 or 13 and a second C subunit amino acid sequence as claimed in claim 12 or 13.
- 40
17. Vascular endothelial cell growth factor according to claim 16 wherein both C subunits have the amino acid sequence claimed in claim 13.
18. Vascular endothelial cell growth factor comprising an A subunit selected from a 189 amino acid form, a 165 amino acid form and a 121 amino acid form, and a C subunit amino acid sequence as claimed in claim 12 or 13.
- 45
19. A tissue repairing pharmaceutical composition comprising a pharmaceutical carrier and an effective tissue repairing amount of the purified vascular endothelial growth factor of any one of claims 14 to 18.
20. The use of the vascular endothelial cell growth factor of any one of claims 14 to 18 for the manufacture of a medicament for promoting tissue repair.
21. The use of the vascular endothelial cell growth factor of any one of claims 14 to 18 for the manufacture of a medicament for stimulating vascular endothelial cell growth.

#### 50 Patentansprüche

1. Gereinigte und isolierte DNA-Sequenz, welche für die C-Untereinheit von Gefäßendothelzell-Wachstumsfaktor kodiert, umfassend:

ATG CCG GTC ATG AGG CTG TTC OCT TGC TTC CTG CAG CTC CTG GCC  
 5 GGG CTG GCG CTG OCT GCT GTG OOC OOC CAG CAG TGG GOC TTG TCT  
 GCT GGG AAC GGC TCG TCA GAG GTG GAA GTG GTA OOC TTC CAG GAA  
 10 GTG TGG GGC OGC AGC TAC TGC OGG GCG CTG GAG AGG CTG GTG GAC  
 GTC GTG TOC GAG TAC COC AGC GAG GTG GAG CAC ATG TTC AGC OCA  
 15 TCC TGT GTC TOC CTG CTG OGC TGC ACC GGC TGC TGC GGC GAT GAG  
 AAT CTG CAC TGT GTG COG GTG GAG ACG GOC AAT GTC AOC ATG CAG  
 CTC CTA AAG ATC OGT TCT GGG GAC OGG OOC TOC TAC GTG GAG CTG  
 20 ACG TTC TCT CAG CAC GTT OGC TGC GAA TGC OGG OCT CTG OGG GAG  
 AAG ATG AAG CCG GAA AGG AGG AGA OOC AAG GGC AGG GGG AAG AGG  
 25 AGG AGA GAG AAG TAG

30 2. Gereinigte und isolierte Gefäßendothelzell-Wachstumsfaktor-C-Untereinheits-DNA-Sequenz, umfassend:

ATG CCG GTC ATG AGG CTG TTC OCT TGC TTC CTG CAG CTC CTG  
 35 GGC GGG CTG GCG CTG OCT GCT GTG OOC OOC CAG CAG TGG GCC  
 TTG TCT GCT GGG AAC GGC TCG TCA GAG GTG GAA GTG GTA CCC  
 TTC CAG GAA GTG TGG GGC OGC AGC TAC TGC OGG GCG CTG GAG  
 40 AGG CTG GTG GAC GTC GTG TOC GAG TAC OOC AGC GAG GTG GAG  
 CAC ATG TTC AGC OCA TOC TGT GTC TOC CTG CTG OGC TGC ACC  
 GGC TGC TGC GGC GAT GAG AAT CTG CAC TGT GTG CCG GTG GAG  
 45 ACG GCC AAT GTC AOC ATG CAG CTC CTA AAG ATC OGT TCT GGG  
 GAC OGG COC TOC TAC GTG GAG CTG ACG TTC TCT CAG CAC GTT  
 OGC TGC GAA TGC OGG OCT CTG OGG GAG AAG ATG AAG CCG GAA  
 AGG AGG AGA COC AAG GGC AGG GGG AAG AGG AGG AGA GAG AAG  
 50 CAG AGA COC ACA GAC TGC CAC CTG TGC GGC GAT GCT GTT CCC  
 CCG AGG TAA. SEQ-ID-Nrn. 28 & 40

55 3. Gefäßendothelzell-Wachstumsfaktor-AC-DNA, umfassend eine A-Untereinheits-DNA-Sequenz wie hier in Figur 4,  
 5 oder 6 dargestellt und eine C-Untereinheits-DNA-Sequenz nach Anspruch 1 oder 2.

4. Gefäßendothelzell-Wachstumsfaktor-BC-DNA, umfassend eine B-Untereinheits-DNA-Sequenz wie hier in Figur 7 oder 8 dargestellt und eine C-Untereinheits-DNA-Sequenz nach Anspruch 1 oder 2.
- 5 5. Gefäßendothelzell-Wachstumsfaktor-AC-DNA, umfassend eine A-Untereinheits-DNA-Sequenz, welche aus der Gruppe, bestehend aus einer DNA-Sequenz, die für eine Form mit 189 Aminosäuren kodiert, einer DNA-Sequenz, die für eine Form mit 165 Aminosäuren kodiert, und einer DNA-Sequenz, die für eine Form mit 121 Aminosäuren kodiert, ausgewählt ist, wobei die A-Untereinheits-DNA funktionsfähig mit einer C-Untereinheits-DNA-Sequenz nach Anspruch 1 oder 2 verknüpft ist.
- 10 6. Gefäßendothelzell-Wachstumsfaktor-BC-DNA, umfassend eine B-Untereinheits-DNA-Sequenz, welche aus der Gruppe, bestehend aus einer DNA-Sequenz, die für eine Form mit 135 Aminosäuren wie hier in Figur 8 dargestellt, mit Ala<sup>24</sup> beginnend und sich bis Leu<sup>158</sup> erstreckend, kodiert, und einer DNA-Sequenz, die für eine Form mit 115 Aminosäuren wie hier in Figur 7 dargestellt, mit Ala<sup>24</sup> beginnend und sich bis Arg<sup>138</sup> erstreckend, kodiert, ausgewählt ist, wobei die B-Untereinheits-DNA-Sequenz funktionsfähig mit einer C-Untereinheits-DNA-Sequenz nach  
15 Anspruch 1 oder 2 verknüpft ist.
7. Homodimere Gefäßendothelzell-Wachstumsfaktor-DNA, umfassend C-Untereinheits-DNA-Sequenzen nach Anspruch 1 oder 2.
- 20 8. Vektor, enthaltend die DNA-Sequenz nach irgendeinem der Ansprüche 2 bis 7.
9. Wirtszelle, transformiert mit dem Vektor nach Anspruch 8, enthaltend die DNA-Sequenz, welche für Gefäßendothelzell-Wachstumsfaktor kodiert.
- 25 10. Verfahren zur Herstellung von Gefäßendothelzell-Wachstumsfaktor, umfassend die Kultivierung der transformierten Wirtszelle nach Anspruch 9 unter Bedingungen, welche für die Expression von Gefäßendothelzell-Wachstumsfaktor und die Gewinnung von Gefäßendothelzell-Wachstumsfaktor geeignet sind.
11. Gefäßendothelzell-Wachstumsfaktor, hergestellt nach dem Verfahren von Anspruch 10.
- 30 12. Gereinigte und isolierte Gefäßendothelzell-Wachstumsfaktor-C-Untereinheits-Aminosäuresequenz, umfassend:

Met Pro Val Met Arg Leu Phe Pro Cys Phe Leu Gln Leu Leu  
 35 Ala Gly Leu Ala Leu Pro Ala Val Pro Pro Gln Gln Trp Ala  
 Leu Ser Ala Gly Asn Gly Ser Ser Glu Val Glu Val Val Pro  
 Phe Gln Glu Val Trp Gly Arg Ser Tyr Cys Arg Ala Leu Glu  
 40 Arg Leu Val Asp Val Val Ser Glu Tyr Pro Ser Glu Val Glu  
 His Met Phe Ser Pro Ser Cys Val Ser Leu Leu Arg Cys Thr  
 Gly Cys Cys Gly Asp Glu Asn Leu His Cys Val Pro Val Glu  
 Thr Ala Asn Val Thr Met Gln Leu Leu Lys Ile Arg Ser Gly  
 45 Asp Arg Pro Ser Tyr Val Glu Leu Thr Phe Ser Gln His Val  
 Arg Cys Glu Cys Arg Pro Leu Arg Glu Lys Met Lys Pro Glu  
 Arg Arg Arg Pro Lys Gly Arg Gly Lys Arg Arg Arg Glu Lys.

50 SEQ-ID-Nr. 41

- 55 13. Gereinigte und isolierte Gefäßendothelzell-Wachstumsfaktor-C-Untereinheits-Aminosäuresequenz, umfassend:

Met Pro Val Met Arg Leu Phe Pro Cys Phe Leu Gln Leu Leu  
 Ala Gly Leu Ala Leu Pro Ala Val Pro Pro Gln Gln Trp Ala  
 5 Leu Ser Ala Gly Asn Gly Ser Ser Glu Val Glu Val Val Pro  
 Phe Gln Glu Val Trp Gly Arg Ser Tyr Cys Arg Ala Leu Glu  
 Arg Leu Val Asp Val Val Ser Glu Tyr Pro Ser Glu Val Glu  
 10 His Met Phe Ser Pro Ser Cys Val Ser Leu Leu Arg Cys Thr  
 Gly Cys Cys Gly Asp Glu Asn Leu His Cys Val Pro Val Glu  
 Thr Ala Asn Val Thr Met Gln Leu Leu Lys Ile Arg Ser Gly  
 15 Asp Arg Pro Ser Tyr Val Glu Leu Thr Phe Ser Gln His Val  
 Arg Cys Glu Cys Arg Pro Leu Arg Glu Lys Met Lys Pro Glu  
 Arg Arg Arg Pro Lys Gly Arg Gly Lys Arg Arg Arg Glu Lys  
 20 Gln Arg Pro Thr Asp Cys His Leu Cys Gly Asp Ala Val Pro  
 Arg Arg. SEQ-ID-Nrn. 29 & 40

- 25
14. Gefäßendothelzell-Wachstumsfaktor AC, umfassend eine A-Untereinheits-Aminosäuresequenz wie hier in Figur 4, 5 oder 6 dargestellt und eine C-Untereinheits-Aminosäuresequenz nach Anspruch 12 oder 13.
- 30
15. Gefäßendothelzell-Wachstumsfaktor BC, umfassend eine B-Untereinheits-Aminosäuresequenz wie hier in Figur 7 oder 8 dargestellt und eine C-Untereinheits-Aminosäuresequenz nach Anspruch 12 oder 13.
16. Gefäßendothelzell-Wachstumsfaktor CC, umfassend eine erste C-Untereinheits-Aminosäuresequenz nach Anspruch 12 oder 13 und eine zweite C-Untereinheits-Aminosäuresequenz nach Anspruch 12 oder 13.
- 35
17. Gefäßendothelzell-Wachstumsfaktor nach Anspruch 16, bei welchem beide C-Untereinheiten die Aminosäuresequenz nach Anspruch 13 aufweisen.
18. Gefäßendothelzell-Wachstumsfaktor, umfassend eine A-Untereinheit, die aus einer Form mit 189 Aminosäuren, einer Form mit 165 Aminosäuren und einer Form mit 121 Aminosäuren ausgewählt ist, und eine C-Untereinheits-Aminosäuresequenz nach Anspruch 12 oder 13.
- 40
19. Gewebe-wiederherstellende pharmazeutische Zusammensetzung, umfassend einen pharmazeutischen Träger und eine wirksame, gewebe-wiederherstellende Menge des gereinigten Gefäßendothelzell-Wachstumsfaktors nach irgendeinem der Ansprüche 14 bis 18.
- 45
20. Verwendung des Gefäßendothelzell-Wachstumsfaktors nach irgendeinem der Ansprüche 14 bis 18 zur Herstellung eines Medikaments zur Förderung der Wiederherstellung von Gewebe.
21. Verwendung des Gefäßendothelzell-Wachstumsfaktors nach irgendeinem der Ansprüche 14 bis 18 zur Herstellung eines Medikaments zur Stimulierung von Gefäßendothelzellwachstum.
- 50

#### Revendications

- 55
1. Séquence d'ADN purifiée et isolée codant pour la sous-unité C du facteur de croissance de cellules endothéliales vasculaires, comprenant :

ATG CCG GTC ATG AGG CTG TTC OCT TGC TTC CTG CAG CTC CTG GCC  
 5 GGG CTG GCG CTG OCT GCT GTG CCC CCC CAG CAG TGG GCC TTG TCT  
 GCT GGG AAC GGC TCG TCA GAG GTG GAA GTG GTA CCC TTC CAG GAA  
 10 GTG TGG GGC CGC AGC TAC TGC OGG GCG CTG GAG AGG CTG GTG GAC  
 GTC GTG TCC GAG TAC CCC AGC GAG GTG GAG CAC ATG TTC AGC OCA  
 TCC TGT GTC TCC CTG CTG CGC TGC ACC GGC TGC TGC GGC GAT GAG  
 15 AAT CTG CAC TGT GTG COG GTG GAG ACG GGC AAT GTC ACC ATG CAG  
 CTC CTA AAG ATC CGT TCT GGG GAC OGG CCC TOC TAC GTG GAG CTG  
 20 ACG TTC TCT CAG CAC GTT CGC TGC GAA TGC OGG OCT CTG OGG GAG  
 AAG ATG AAG OCG GAA AGG AGG AGA CCC AAG GGC AGG GGG AAG AGG  
 AGG AGA GAG AAG TAG

2. Séquence d'ADN purifiée et isolée de la sous-unité C du facteur de croissance de cellules endothéliales vasculaires, comprenant :

30 ATG CCG GTC ATG AGG CTG TTC OCT TGC TTC CTG CAG CTC CTG  
 GCC GGG CTG GCG CTG OCT GCT GTG CCC CCC CAG CAG TGG GCC  
 35 TTG TCT GCT GGG AAC GGC TCG TCA GAG GTG GAA GTG GTA CCC  
 TTC CAG GAA GTG TGG GGC CGC AGC TAC TGC OGG GCG CTG GAG  
 AGG CTG GTG GAC GTC GTG TOC CAG TAC CCC AGC GAG GTG GAG  
 40 CAC ATG TTC AGC OCA TOC TGT GTC TOC CTG CTG CGC TGC ACC  
 GGC TGC TGC GGC GAT GAG AAT CTG CAC TGT GTG OCG GTG GAG  
 ACG GGC AAT GTC AOC ATG CAG CTC CTA AAG ATC CGT TCT GGG  
 GAC CGG CCC TOC TAC GTG GAG CTG ACG TTC TCT CAG CAC GTT  
 45 CGC TGC GAA TGC CGG CCT CTG OGG GAG AAG ATG AAG CCC GAA  
 AAG AGG AGA CCC AAG GGC AGG GGG AAG AGG AGG AGA GAG AAG  
 CAG AGA CCC ACA GAC TGC CAC CTG TGC GGC GAT GCT GTT CCC  
 50 OGG AGG TAA. SEQ ID NOS: 23 & 40

3. ADN du facteur AC de croissance de cellules endothéliales vasculaires comprenant une séquence d'ADN de la sous-unité A telle que représentée sur les figures 4, 5 ou 6 annexées et une séquence d'ADN de la sous-unité C selon la revendication 1 ou 2.

4. ADN du facteur BC de croissance de cellules endothéliales vasculaires comprenant une séquence d'ADN de la

sous-unité B telle que représentée sur les figures 7 ou 8 annexées et une séquence d'ADN de la sous-unité C selon la revendication 1 ou 2.

- 5 5. ADN du facteur AC de croissance de cellules endothéliales vasculaires comprenant une séquence d'ADN de la sous-unité A sélectionnée parmi le groupe constitué d'une séquence d'ADN codant pour une forme à 189 acides aminés, d'une séquence d'ADN codant pour une forme à 165 acides aminés et d'une séquence d'ADN codant pour une forme à 121 acides aminés, ledit ADN de la sous-unité A étant attaché de manière opérationnelle à une séquence d'ADN de la sous-unité C selon la revendication 1 ou 2.
- 10 6. ADN du facteur BC de croissance de cellules endothéliales vasculaires comprenant une séquence d'ADN de la sous-unité B sélectionnée parmi le groupe constitué d'une séquence d'ADN codant pour une forme à 135 acides aminés telle que représentée sur la figure 8 annexée, commençant à Ala<sup>24</sup> et se poursuivant jusqu'à Leu<sup>158</sup>, et d'une séquence d'ADN codant pour une forme à 115 acides aminés telle que représentée sur la figure 7 annexée, commençant à Ala<sup>24</sup> et se poursuivant jusqu'à Arg<sup>138</sup>, ladite séquence d'ADN de la sous-unité B étant attachée de manière opérationnelle à une séquence d'ADN de la sous-unité C selon la revendication 1 ou 2.
- 15 7. ADN du facteur de croissance de cellules endothéliales vasculaires homodimère, comprenant des séquences d'ADN de la sous-unité C selon la revendication 1 ou 2.
- 20 8. Vecteur contenant la séquence d'ADN selon l'une quelconque des revendications 2 à 7.
9. Cellule hôte transformée par le vecteur selon la revendication 8, contenant la séquence d'ADN codant pour le facteur de croissance de cellules endothéliales vasculaires.
- 25 10. Procédé de préparation d'un facteur de croissance de cellules endothéliales vasculaires comprenant les étapes consistant à mettre en culture la cellule hôte transformée selon la revendication 9 dans des conditions adaptées à l'expression du facteur de croissance de cellules endothéliales vasculaires et récupérer le facteur de croissance de cellules endothéliales vasculaires.
- 30 11. Facteur de croissance de cellules endothéliales vasculaires produit par le procédé de la revendication 10.
12. Séquence d'acides aminés purifiée et isolée de la sous-unité C du facteur de croissance de cellules endothéliales vasculaires, comprenant :

35 Met Pro Val Met Arg Leu Phe Pro Cys Phe Leu Gln Leu Leu  
 Ala Gly Leu Ala Leu Pro Ala Val Pro Pro Gln Gln Trp Ala  
 Leu Ser Ala Gly Asn Gly Ser Ser Glu Val Glu Val Val Pro  
 40 Phe Gln Glu Val Trp Gly Arg Ser Tyr Cys Arg Ala Leu Glu  
 Arg Leu Val Asp Val Val Ser Glu Tyr Pro Ser Glu Val Glu  
 His Met Phe Ser Pro Ser Cys Val Ser Leu Leu Arg Cys Thr  
 45 Gly Cys Cys Gly Asp Glu Asn Leu His Cys Val Pro Val Glu  
 Thr Ala Asn Val Thr Met Gln Leu Leu Lys Ile Arg Ser Gly  
 Asp Arg Pro Ser Tyr Val Glu Leu Thr Phe Ser Gln His Val  
 Arg Cys Glu Cys Arg Pro Leu Arg Glu Lys Met Lys Pro Glu  
 50 Arg Arg Arg Pro Lys Gly Arg Gly Lys Arg Arg Arg Glu Lys.

SEQ ID NO:41

55

13. Séquence d'acides aminés purifiée et isolée de la sous-unité C du facteur de croissance de cellules endothéliales vasculaires, comprenant :

Met Pro Val Met Arg Leu Phe Pro Cys Phe Leu Gln Leu Leu  
 Ala Gly Leu Ala Leu Pro Ala Val Pro Pro Gln Gln Trp Ala  
 5 Leu Ser Ala Gly Asn Gly Ser Ser Glu Val Glu Val Val Pro  
 Phe Gln Glu Val Trp Gly Arg Ser Tyr Cys Arg Ala Leu Glu  
 Arg Leu Val Asp Val Val Ser Glu Tyr Pro Ser Glu Val Glu  
 10 His Met Phe Ser Pro Ser Cys Val Ser Leu Leu Arg Cys Thr  
 Gly Cys Cys Gly Asp Glu Asn Leu His Cys Val Pro Val Glu  
 Thr Ala Asn Val Thr Met Gln Leu Leu Lys Ile Arg Ser Gly  
 Asp Arg Pro Ser Tyr Val Glu Leu Thr Phe Ser Gln His Val  
 15 Arg Cys Glu Cys Arg Pro Leu Arg Glu Lys Met Lys Pro Glu  
 Arg Arg Arg Pro Lys Gly Arg Gly Lys Arg Arg Arg Glu Lys  
 Gln Arg Pro Thr Asp Cys His Leu Cys Gly Asp Ala Val Pro  
 20 Arg Arg. SEQ ID NOS: 29 & 40

25 14. Facteur AC de croissance de cellules endothéliales vasculaires, comprenant une séquence d'acides aminés de la sous-unité A telle que représentée sur les figures 4, 5 ou 6 annexées et une séquence d'acides aminés de la sous-unité C selon la revendication 12 ou 13.

30 15. Facteur BC de croissance de cellules endothéliales vasculaires comprenant une séquence d'acides aminés de la sous-unité B telle que représentée sur les figures 7 ou 8 annexées et une séquence d'acides aminés de la sous-unité C selon la revendication 12 ou 13.

35 16. Facteur CC de croissance de cellules endothéliales vasculaires comprenant une première séquence d'acides aminés de la sous-unité C selon la revendication 12 ou 13 et une seconde séquence d'acides aminés de la sous-unité C selon la revendication 12 ou 13.

40 17. Facteur de croissance de cellules endothéliales vasculaires selon la revendication 16, dans lequel les deux sous-unités C ont la séquence d'acides aminés de la revendication 13.

18. Facteur de croissance de cellules endothéliales vasculaires comprenant une sous-unité A sélectionnée parmi une forme à 189 acides aminés, une forme à 165 acides aminés et une forme à 121 acides aminés, et une séquence d'acides aminés de la sous-unité C selon la revendication 12 ou 13.

45 19. Composition pharmaceutique réparatrice de tissu comprenant un véhicule pharmaceutique et une quantité efficace pour la réparation de tissu de facteur de croissance endothéliale vasculaire purifié selon l'une des revendications 14 à 18.

50 20. Utilisation du facteur de croissance de cellules endothéliales vasculaires selon l'une des revendications 14 à 18 pour la préparation d'un médicament destiné à favoriser la réparation de tissu.

21. Utilisation du facteur de croissance de cellules endothéliales vasculaires selon l'une des revendications 14 à 18 pour la préparation d'un médicament destiné à stimuler la croissance des cellules endothéliales vasculaires.



FIG. 1




FIG. 1 (CONT.)




FIG. / (CONT.)




FIG. / (CONT.)




FIG. 1 (CONT.)




FIG. 2




FIG. 2 (CONT.)

p4238	
TTC CAG GAG TAC CCC GAT GAG ATA GAG TAT ATC TTC ARG CCG TCC TGT GTG CCC CTA ATG	80
PHE-GLN-GLU-TYR-PRO-ASP-GLU-ILE-GLU-TYR-ILE-PHE-LYS-PRO-SER-CYS-VAL-PRO-LEU-MET-	90
→ → → → → → → →	
L46	
p4238	
CGG TGT GCG GGC TGC TGC AAT GAT GAA GCC CTG GAG TGC GTG CCC ACG TCG GAG AGC AAC	100
ARG-CYS-ALA-GLY-CYS-CYS-ASN-ASP-GLU-ALA-LEU-GLU-CYS-VAL-PRO-THR-SER-GLU-SER-ASN-	110
→ → → → → → → →	
L46	
p4238	
GTC ACT ATG CAG ATC CGG ATC AAA CCT CAC CAA AGC CAG CAC ATA GGA GAG ATG AGC	120
VAL-THR-MET-GLN-ILE-MET-ARG-ILE-LYS-PRO-HIS-GLN-SER-GLN-HIS-ILE-GLY-GLU-MET-SER-	130
→ → → → → → → →	
L46	

FIG. 2 (CONT.)






FIG. 2 (CONT.)




FIG. 3




FIG. 3 (CONT.)





FIG. 3 (CONT.)



CTG TGA  
 158  
 LEU •

FIG. 4

ACCA	ATG	AAC	TTT	CTG	CTC	TCT	TGG	GTG	CAC	TGG	ACC	CTG	GCT	TTA	CTG	49
Met	Asn	Phe	Leu	Leu	Ser	Trp	Val	His	Trp	Thr	Leu	Ala	Leu	Leu		
				5					10					15		
CTG	TAC	CTC	CAC	CAT	GCC	AAG	TGG	TOC	CAG	GCT	GCA	CCC	ACG	ACA	94	
Leu	Tyr	Leu	His	His	Ala	Lys	Trp	Ser	Gln	Ala	Ala	Pro	Thr	Thr		
				20					25					30		
GAA	GGG	GAG	CAG	AAA	GCC	CAT	GAA	GTG	GTG	AAG	TTC	ATG	GAC	GTC	139	
Glu	Gly	Glu	Gln	Lys	Ala	His	Glu	Val	Val	Lys	Phe	Met	Asp	Val		
				35					40					45		
TAC	CAG	CGC	AGC	TAT	TGC	OGT	CCG	ATT	GAG	ACC	CTG	GTG	GAC	ATC	184	
Tyr	Gln	Arg	Ser	Tyr	Cys	Arg	Pro	Ile	Glu	Thr	Leu	Val	Asp	Ile		
				50					55					60		
TTC	CAG	GAG	TAC	CCC	GAT	GAG	ATA	GAG	TAT	ATC	TTC	AAG	CCG	TOC	229	
Phe	Gln	Glu	Tyr	Pro	Asp	Glu	Ile	Glu	Tyr	Ile	Phe	Lys	Pro	Ser		
				65					70					75		
TGT	GTG	CCC	CTA	ATG	CGG	TGT	GCG	GGC	TGC	TGC	AAT	GAT	GAA	GCC	274	
Cys	Val	Pro	Leu	Met	Arg	Cys	Ala	Gly	Cys	Cys	Asn	Asp	Glu	Ala		
				80					85					90		
CTG	GAG	TGC	GTG	CCC	ACG	TOG	GAG	AGC	AAC	GTC	ACT	ATG	CAG	ATC	319	
Leu	Glu	Cys	Val	Pro	Thr	Ser	Glu	Ser	Asn	Val	Thr	Met	Gln	Ile		
				95					100					105		
ATG	CGG	ATC	AAA	OCT	CAC	CAA	AGC	CAG	CAC	ATA	GGA	GAG	ATG	AGC	364	
Met	Arg	Ile	Lys	Pro	His	Gln	Ser	Gln	His	Ile	Gly	Glu	Met	Ser		
				110					115					120		
TTC	CTG	CAG	CAT	AGC	AGA	TGT	GAA	TGC	AGA	CCA	AAG	AAA	GAT	AGA	409	
Phe	Leu	Gln	His	Ser	Arg	Cys	Glu	Cys	Arg	Pro	Lys	Lys	Asp	Arg		
				125					130					135		
ACA	AAG	CCA	GAA	AAA	TGT	GAC	AAG	CCA	AGG	OGG	TGA				445	
Thr	Lys	Pro	Glu	Lys	Cys	Asp	Lys	Pro	Arg	Arg						
				140					145							

FIG. 5

AAC	ATG	AAC	TTT	CTG	CTC	TCT	TGG	GTG	CAC	TGG	AOC	CTG	GCT	TTA	CTG	49
Met	Asn	Phe	Leu	Leu	Ser	Trp	Val	His	Trp	Thr	Leu	Ala	Leu	Leu		
				5						10					15	
CTG	TAC	CTC	CAC	CAT	GOC	AAG	TGG	TOC	CAG	GCT	GCA	OCG	ACG	ACA		94
Leu	Tyr	Leu	His	His	Ala	Lys	Trp	Ser	Gln	Ala	Ala	Pro	Thr	Thr		
				20					25					30		
GAA	GGG	GAG	CAG	AAA	GOC	CAT	GAA	GTG	GTG	AAG	TTC	ATG	GAC	GTC		139
Glu	Gly	Glu	Gln	Lys	Ala	His	Glu	Val	Val	Lys	Phe	Met	Asp	Val		
				35					40					45		
TAC	CAG	CGC	AGC	TAT	TGC	CGT	CCG	ATT	GAG	AOC	CTG	GTG	GAC	ATC		184
Tyr	Gln	Arg	Ser	Tyr	Cys	Arg	Pro	Ile	Glu	Thr	Leu	Val	Asp	Ile		
				50					55					60		
TTC	CAG	GAG	TAC	CCC	GAT	GAG	ATA	GAG	TAT	ATC	TTC	AAG	CCG	TOC		229
Phe	Gln	Glu	Tyr	Pro	Asp	Glu	Ile	Glu	Tyr	Ile	Phe	Lys	Pro	Ser		
				65					70					75		
TGT	GTG	CCC	CTA	ATG	CGG	TGT	GCG	GGC	TGC	TGC	AAT	GAT	GAA	GOC		274
Cys	Val	Pro	Leu	Met	Arg	Cys	Ala	Gly	Cys	Cys	Asn	Asp	Glu	Ala		
				80					85					90		
CTG	GAG	TGC	GTG	CCC	ACG	TOG	GAG	AGC	AAC	GTC	ACT	ATG	CAG	ATC		319
Leu	Glu	Cys	Val	Pro	Thr	Ser	Glu	Ser	Asn	Val	Thr	Met	Gln	Ile		
				95					100					105		
ATG	CGG	ATC	AAA	OCT	CAC	CAA	AGC	CAG	CAC	ATA	GGA	GAG	ATG	AGC		364
Met	Arg	Ile	Lys	Pro	His	Gln	Ser	Gln	His	Ile	Gly	Glu	Met	Ser		
				110					115					120		
TTC	CTG	CAG	CAT	AGC	AGA	TGT	GAA	TGC	AGA	CCA	AAG	AAA	GAT	AGA		409
Phe	Leu	Gln	His	Ser	Arg	Cys	Glu	Cys	Arg	Pro	Lys	Lys	Asp	Arg		
				125					130					135		
ACA	AAG	CCA	GAA	AAT	CAC	TGT	GAG	OCT	TGT	TCA	GAG	CGG	AGA	AAG		454
Thr	Lys	Pro	Glu	Asn	His	Cys	Glu	Pro	Cys	Ser	Glu	Arg	Arg	Lys		
				140					145					150		
CAT	TTG	TTT	GTC	CAA	GAT	CCG	CAG	ACG	TGT	AAA	TGT	TOC	TGC	AAA		499
His	Leu	Phe	Val	Gln	Asp	Pro	Gln	Thr	Cys	Lys	Cys	Ser	Cys	Lys		
				155					160					165		
AAC	ACA	GAC	TOG	CGT	TGC	AA	GCG	AGG	CAG	CTT	GAG	TTA	AAC	GAA		544
Asn	Thr	Asp	Ser	Arg	Cys	Lys	Ala	Arg	Gln	Leu	Glu	Leu	Asn	Glu		
				170					175					180		
CGT	ACT	TGC	AGA	TGT	GAC	AAG	CCA	AGG	CGG	TGA						577
Arg	Thr	Cys	Arg	Cys	Asp	Lys	Pro	Arg	Arg							
				185					190							

FIG. 6A

AACC ATG AAC TTT CTG CTC TCT TGG GTG CAC TGG ACC CTG GCT TTA CTG	49
Met Asn Phe Leu Leu Ser Trp Val His Trp Thr Leu Ala Leu Leu	
5 10 15	
CTG TAC CTC CAC CAT GOC AAG TGG TOC CAG GCT GCA CCC ACG ACA	94
Leu Tyr Leu His His Ala Lys Trp Ser Gln Ala Ala Pro Thr Thr	
20 25 30	
GAA GGG GAG CAG AAA GOC CAT GAA GTG GTG AAG TTC ATG GAC GTC	139
Glu Gly Glu Gln Lys Ala His Glu Val Val Lys Phe Met Asp Val	
35 40 45	
TAC CAG CGC AGC TAT TGC CGT CCG ATT GAG ACC CTG GTG GAC ATC	184
Tyr Gln Arg Ser Tyr Cys Arg Pro Ile Glu Thr Leu Val Asp Ile	
50 55 60	
TTC CAG GAG TAC CCC GAT GAG ATA GAG TAT ATC TTC AAG CCG TOC	229
Phe Gln Glu Tyr Pro Asp Glu Ile Glu Tyr Ile Phe Lys Pro Ser	
65 70 75	
TGT GTG CCC CTA ATG CGG TGT GCG GGC TGC TGC AAT GAT GAA GOC	274
Cys Val Pro Leu Met Arg Cys Ala Gly Cys Cys Asn Asp Glu Ala	
80 85 90	
CTG GAG TGC GTG CCC ACG TOG GAG AGC AAC GTC ACT ATG CAG ATC	319
Leu Glu Cys Val Pro Thr Ser Glu Ser Asn Val Thr Met Gln Ile	
95 100 105	
ATG CGG ATC AAA OCT CAC CAA AGC CAG CAC ATA GGA GAG ATG AGC	364
Met Arg Ile Lys Pro His Gln Ser Gln His Ile Gly Glu Met Ser	
110 115 120	
TTC CTG CAG CAT AGC AGA TGT GAA TGC AGA CCA AAG AAA GAT AGA	409
Phe Leu Gln His Ser Arg Cys Glu Cys Arg Pro Lys Lys Asp Arg	
125 130 135	
ACA AAG CCA GAA AAA AAA TCA GTT CGA GGA AAG GGA AAG GGT CAA	454
Thr Lys Pro Glu Lys Lys Ser Val Arg Gly Lys Gly Lys Gly Gln	
140 145 150	
AAA CGA AAG CGC AAG AAA TOC CGG TTT AAA TOC TGG AGC GTT CAC	499
Lys Arg Lys Arg Lys Lys Ser Arg Phe Lys Ser Trp Ser Val His	
155 160 165	

FIG 6B

TGT	GAG	OCT	TGT	TCA	GAG	CGG	AGA	AAG	CAT	TTG	TTT	GTC	CAA	GAT	544
Cys	Glu	Pro	Cys	Ser	Glu	Arg	Arg	Lys	His	Leu	Phe	Val	Gln	Asp	
				170					175					180	
CCG	CAG	ACG	TGT	AAA	TGT	TOC	TGC	AAA	AAC	ACA	GAC	TOG	CGT	TGC	589
Pro	Gln	Thr	Cys	Lys	Cys	Ser	Cys	Lys	Asn	Thr	Asp	Ser	Arg	Cys	
				185					190					195	
AAG	GCG	AGG	CAG	CTT	GAG	TTA	AAC	GAA	CGT	ACT	TGC	AGA	TGT	GAC	634
Lys	Ala	Arg	Gln	Leu	Glu	Leu	Asn	Glu	Arg	Thr	Cys	Arg	Cys	Asp	
				200					205					210	
AAG	CCA	AGG	CGG	TGA											645
Lys	Pro	Arg	Arg												



FIG. 7

ATG CTG GCC ATG AAG CTG TTC ACT TGC TTC TTG CAG GTC CTA GCT	45
Met Leu Ala Met Lys Leu Phe Thr Cys Phe Leu Gln Val Leu Ala	
5 10 15	
GGG TTG GCT GTG CAC TOC CAG GGG GGC CTG TCT GCT GGG AAC AAC	90
Gly Leu Ala Val His Ser Gln Gly Ala Leu Ser Ala Gly Asn Asn	
20 25 30	
TCA ACA GAA ATG GAA GTG GTG OCT TTC AAT GAA GTG TGG GGC CGC	135
Ser Thr Glu Met Glu Val Val Pro Phe Asn Glu Val Trp Gly Arg	
35 40 45	
AGC TAC TGC CGG CCA ATG GAG AAG CTG GTG TAC ATT GCA GAT GAA	180
Ser Tyr Cys Arg Pro Met Glu Lys Leu Val Tyr Ile Ala Asp Glu	
50 55 60	
CAC OCT AAT GAA GTG TCT CAT ATA TTC AGT CCG TCA TGT GTC CTT	225
His Pro Asn Glu Val Ser His Ile Phe Ser Pro Ser Cys Val Leu	
65 70 75	
CTG AGT CGC TGT AGT GGC TGC TGT GGT GAC GAG GGT CTG CAC TGT	270
Leu Ser Arg Cys Ser Gly Cys Cys Gly Asp Glu Gly Leu His Cys	
80 85 90	
GTG GCG CTA AAG ACA GCC AAC ATC ACT ATG CAG ATC TTA AAG ATT	315
Val Ala Leu Lys Thr Ala Asn Ile Thr Met Gln Ile Leu Lys Ile	
95 100 105	
CCC CCC AAT CGG GAT CCA CAT TOC TAC GTG GAG ATG ACA TTC TCT	360
Pro Pro Asn Arg Asp Pro His Ser Tyr Val Glu Met Thr Phe Ser	
110 115 120	
CAG GAT GTA CTC TGC GAA TGC AGG OCT ATT CTG GAG ACG ACA AAG	405
Gln Asp Val Leu Cys Glu Cys Arg Pro Ile Leu Glu Thr Thr Lys	
125 130 135	
GCA GAA AGG TAA	417
Ala Glu Arg	

FIG. 8

ATG CTG GCC ATG AAG CTG TTC ACT TGC TTC TTG CAG GTC CTA GCT	45
Met Leu Ala Met Lys Leu Phe Thr Cys Phe Leu Gln Val Leu Ala	
5 10 15	
GGG TTG GCT GTG CAC TOC CAG GGG GOC CTG TCT GCT GGG AAC AAC	90
Gly Leu Ala Val His Ser Gln Gly Ala Leu Ser Ala Gly Asn Asn	
20 25 30	
TCA ACA GAA ATG GAA GTG GTG OCT TTC AAT GAA GTG TGG GGC GGC	135
Ser Thr Glu Met Glu Val Val Pro Phe Asn Glu Val Trp Gly Arg	
35 40 45	
AGC TAC TGC CCG CCA ATG GAG AAG CTG GTG TAC ATT GCA GAT GAA	180
Ser Tyr Cys Arg Pro Met Glu Lys Leu Val Tyr Ile Ala Asp Glu	
50 55 60	
CAC CCT AAT GAA GTG TCT CAT ATA TTC AGT CCG TCA TGT GTC CTT	225
His Pro Asn Glu Val Ser His Ile Phe Ser Pro Ser Cys Val Leu	
65 70 75	
CTG AGT CCG TGT AGT GGC TGC TGT GGT GAC GAG GGT CTG CAC TGT	270
Leu Ser Arg Cys Ser Gly Cys Cys Gly Asp Glu Gly Leu His Cys	
80 85 90	
GTG GCG CTA AAG ACA GCC AAC ATC ACT ATG CAG ATC TTA AAG ATT	315
Val Ala Leu Lys Thr Ala Asn Ile Thr Met Gln Ile Leu Lys Ile	
95 100 105	
CCC CCC AAT CCG GAT CCA CAT TOC TAC GTG GAG ATG ACA TTC TCT	360
Pro Pro Asn Arg Asp Pro His Ser Tyr Val Glu Met Thr Phe Ser	
110 115 120	
CAG GAT GTA CTC TGC GAA TGC AGG OCT ATT CTG GAG ACG ACA AAG	405
Gln Asp Val Leu Cys Glu Cys Arg Pro Ile Leu Glu Thr Thr Lys	
125 130 135	
GCA GAA AGG AGG AAA ACC AAG GGG AAG AGG AAG CAA AGC AAA ACC	450
Ala Glu Arg Arg Lys Thr Lys Gly Lys Arg Lys Gln Ser Lys Thr	
140 145 150	
OCA CAG ACT GAG GAA CCC CAC CTG TGA	477
Pro Gln Thr Glu Glu Pro His Leu	
155	

FIG. 9

ATG CCG GTC ATG AGG CTG TTC OCT TGC TTC CTG CAG CTC CTG GGC	45
Met Pro Val Met Arg Leu Phe Pro Cys Phe Leu Gln Leu Leu Ala	
5 10 15	
GGG CTG GCG CTG CCT GCT GTG CCC CCC CAG CAG TGG GGC TTG TCT	90
Gly Leu Ala Leu Pro Ala Val Pro Pro Gln Gln Trp Ala Leu Ser	
20 25 30	
GCT GGG AAC GGC TCG TCA GAG GTG GAA GTG GTA CCC TTC CAG GAA	135
Ala Gly Asn Gly Ser Ser Glu Val Glu Val Val Pro Phe Gln Glu	
35 40 45	
GTG TGG GGC CGC AGC TAC TGC CCG GCG CTG GAG AGG CTG GTG GAC	180
Val Trp Gly Arg Ser Tyr Cys Arg Ala Leu Glu Arg Leu Val Asp	
50 55 60	
GTC GTG TCC GAG TAC CCC AGC GAG GTG GAG CAC ATG TTC AGC CCA	225
Val Val Ser Glu Tyr Pro Ser Glu Val Glu His Met Phe Ser Pro	
65 70 75	
TCC TGT GTC TCC CTG CTG CGC TGC ACC GGC TGC TGC GGC GAT GAG	270
Ser Cys Val Ser Leu Leu Arg Cys Thr Gly Cys Cys Gly Asp Glu	
80 85 90	
AAT CTG CAC TGT GTG CCG GTG GAG ACG GCC AAT GTC ACC ATG CAG	315
Asn Leu His Cys Val Pro Val Glu Thr Ala Asn Val Thr Met Gln	
95 100 105	
CTC CTA AAG ATC CGT TCT GGG GAC CCG CCC TCC TAC GTG GAG CTG	360
Leu Leu Lys Ile Arg Ser Gly Asp Arg Pro Ser Tyr Val Glu Leu	
110 115 120	
ACG TTC TCT CAG CAC GTT CGC TGC GAA TGC CCG CCT CTG CCG GAG	405
Thr Phe Ser Gln His Val Arg Cys Glu Cys Arg Pro Leu Arg Glu	
125 130 135	
AAG ATG AAG CCG GAA AGG AGG AGA CCC AAG GGC AGG GGG AAG AGG	450
Lys Met Lys Pro Glu Arg Arg Arg Pro Lys Gly Arg Gly Lys Arg	
140 145 150	
AGG AGA GAG AAG TAG	465
Arg Arg Glu Lys	